



NEW EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the opposition decision:
03.11.1999 Bulletin 1999/44

(51) Int Cl.⁶: C12N 15/13, C12N 15/70,
C12N 15/85, C12N 1/21,
C12N 5/10

(45) Mention of the grant of the patent:
02.03.1994 Bulletin 1994/09

(86) International application number:
PCT/US88/02514

(21) Application number: 88907510.7

(87) International publication number:
WO 89/00999 (09.02.1989 Gazette 1989/04)

(22) Date of filing: 25.07.1988

(54) MODULAR ASSEMBLY OF ANTIBODY GENES, ANTIBODIES PREPARED THEREBY AND USE
MODULÄRE ZUSAMMENBAU VON ANTIKÖRPERGENEN, DADURCH HERGESTELLTE
ANTIKÖRPER UND DEREN ANWENDUNG
ENSEMBLE MODULAIRE DE GENES D'ANTICORPS, ANTICORPS AINSI PREPARES ET
UTILISATION

(84) Designated Contracting States:
AT BE CH DE FR GB IT LI LU NL SE

(30) Priority: 24.07.1987 US 77528

(43) Date of publication of application:
13.06.1990 Bulletin 1990/24

(60) Divisional application: 93100041.8 / O 550 400

(73) Proprietor: XOMA Corporation
Berkeley California 94710-2737 (US)

(72) Inventors:
• ROBINSON, Randy, R.
Los Angeles, CA 90045 (US)
• LIU, Alvin, Y.
Oceanside, CA 92056 (US)
• HORWITZ, Arnold, H.
Los Angeles, CA 90045 (US)
• WALL, Randolph
Sherman Oaks, CA 91423 (US)
• BETTER, Marc
Los Angeles, CA 90046 (US)

(74) Representative: Rlter, Stephen David et al
Mathys & Squire
100 Grays Inn Road
London WC1X 8AL (GB)

(56) References cited:

EP-A- 0 125 023	EP-A- 0 171 496
EP-A- 0 173 494	EP-A- 0 184 187
EP-A- 0 234 592	EP-A1- 0 120 694
EP-A1- 0 126 338	EP-A1- 0 154 539
WO-A-86/01533	WO-A-87/02671
GB-A- 2 137 631	

- Journal of Immunology (Baltimore) Volume 137 Number 3, issued 01 August 1986, SAHAGAN et al. "A genetically engineered murine human chimeric antibody retains specificity for human tumor-associated antigen See pages 1066-1074.
- Nature (London) Volume 309, 24 May 1984, SHARON et al. "Expression or VHCK chimeric protein in mouse myeloma cells" See pages 364-367.
- Nature (London) Volume 312, issued 13 December 1984, Boullianne et al. "Production of functional chimeric mouse/human antibody" see pages 643-646.
- Proceedings of the National Academy of Science USA (Washington, D.C.) Volume 81, issued June 1984, Cabilly et al. "Generation of antibody activity from immunoglobulin chains produced in Escherichia coli" see page 3273-3277.
- Biotechniques (Natick, MA) Volume 4 issued May-June 1986, Ol et al "Chimeric Antibodies" see pages 216-220.
- Nature (London) Volume 314, issued 04 April 1985, TAKEDA et al. "Construction of chimeric processed immunoglobulin genes containing mouse variable and human constant region sequences" see pages 452-454
- Proceedings of the National Academy of Sciences, USA (Washington, D.C.) Volume 81 Issued November 1984, MORRISON et al., "Chimeric human antibody molecules : mouse antigen-binding domains with human constant region domains" see pages 6851-6855.
- Science, (Washington, DC) Volume 229 Issued 20 September 1985, MORRISON "Transfectomas provide novel chimeric antibodies" see pages 1205-1207.

- Science, (Washington, D.C.) Volume 240 Issued 20 May 1988, BETTER et al. pages 1041-1043
- Nucleic Acids Research, (Oxford) Volume 12 Issued December 1984, KRAMER et al. "The gapped duplex DNA approach to oligonucleotide-directed mutation construction". See pages 9441-9455.
- BioTechnology (Martinsville, NJ) Volume 4 Issued November 1986, HSIUNG et al. "High-level expression efficient secretion and folding of human growth hormone in *Escherichia coli*" see pages 991-995 particularly p. 992 and figure 3.
- Nature (London) Volume 312 Issued 13 December 1984, NEUBERGER et al. "Recombinant antibodies possessing novel effector functions" see pages 604-608, particularly pages 605-606 and figures 1 and 2.
- Nature (London) Volume 314 Issued 21 March 1985, NEUBERGER et al. "A hapten-specific chimaeric IgE antibody with human physiological effector function" see pages 268-270.
- Journal of Immunology (Baltimore) Volume 135 Number 5 Issued November 1985, TAN et al. "A human-mouse chimeric immunoglobulin gene with a human variable region is expressed in mouse myeloma cells" see pages 8564-8567.
- Gene, (Amsterdam) Volume 43. Issued 25 August 1986, Williams et al. "Production of antibody-tagged enzymes by myeloma cells: application to DNA polymerase I Klenow fragment" see pages 319-324.
- Nature, (London) Volume 314 04 April 1985, Wood et al. "The synthesis and in vivo assembly of functional antibodies in yeast" see pages 446-448.
- Proc. Natl. Sci. USA 80 (1983), pp 6351-6355.
- Proc. Natl. Acad. Sci. USA 84 (1987), pp. 3439-3443.
- The EMBO Journal, 44, No. 1, pp. 519-526, 1985; McCarthy et al
- Amino Acid and Nucleotide Biosynthesis, pp. 284-287; Jones E.W.
- PNAS, 83, pp. 8508-8510; Schoner et al
- Methods in Enzymology, Goeddel, ed. Academic Press, 1990, pp. 166-187; Stader et al
- Protein Engineering, 2(3), 1988, pp. 169-170; Wetzel
- BioTechn., 10, 1992, pp. 163-167; Carter et al
- Nucl. Acid Res., 12, 1984, pp.3791-3806; Boss et al
- Science, 240, 1988, pp. 1038-1041; Skerra et al
- PNAS, 52, 1964, pp. 1099-1106; Haber et al
- Adv. Immunol., 44, 1989, pp. 65-92; Morrison & Oi

Remarks:

Divisional application 93100041.6 filed on 25/07/88.

Description

Field of the Invention

[0001] This invention relates to recombinant DNA methods of preparing immunoglobulins, genetic sequences coding therefor, as well as methods of obtaining such sequences.

Background Art

[0002] The application of cell-to-cell fusion for the production of monoclonal antibodies by Kohler and Milstein (Nature (London), 256: 495, 1975) has spawned a revolution in biology equal in impact to the invention of recombinant DNA cloning. Hybridoma-produced monoclonal antibodies are already widely used in clinical diagnoses and basic scientific studies. Applications of human B cell hybridoma-produced monoclonal antibodies hold great promise for the clinical treatment of cancer, viral and microbial infections, B cell immunodeficiencies with diminished antibody production, and other diseases and disorders of the immune system.

[0003] Unfortunately, yields of monoclonal antibodies from human hybridoma cell lines are relatively low (1 ug/ml in human x human compared to 100 ug/ml in mouse hybridomas), and production costs are high for antibodies made in large scale human tissue culture. Mouse x mouse hybridomas, on the other hand, are useful because they produce abundant amounts of protein, and these cell lines are more stable than the human lines. However, repeated injections of "foreign" antibodies, such as a mouse antibody, in humans, can lead to harmful hypersensitivity reactions.

[0004] There has therefore been recent exploration of the possibility of producing antibodies having the advantages of monoclonals from mouse-mouse hybridomas, yet the species specific properties of human monoclonal antibodies.

[0005] Another problem faced by immunologists is that most human monoclonal antibodies (i.e., antibodies having human recognition properties) obtained in cell culture are of the IgM type. When it is desirable to obtain human monoclonals of the IgG type, however, it has been necessary to use such techniques as cell sorting, to separate the few cells which have switched to producing antibodies of the IgG or other type from the majority producing antibodies of the IgM type. A need therefore exists for a more ready method of switching antibody classes, for any given antibody of a predetermined or desired antigenic specificity.

[0006] The present invention bridges both the hybridoma and monoclonal antibody technologies and provides a quick and efficient method, as well as products derived therefrom, for the improved production of chimeric human/non-human antibodies, or of "class switched" antibodies.

[0007] Approaches to the problem of producing chimeric antibodies have been published by various authors.

[0008] Morrison, S. L. et al., Proc. Natl. Acad. Sci., USA, 81: 6851-6855 (November 1984), describe the production of a mouse-human antibody molecule of defined antigen binding specificity, produced by joining the variable region of genes of a mouse antibody-producing myeloma cell line with known antigen binding specificity to human immunoglobulin constant region genes using recombinant DNA techniques. Chimeric genes were constructed, wherein the heavy chain variable region exon from the myeloma cell line S107 well joined to human IgG1 or IgG2 heavy chain constant region exons, and the light chain variable region exon from the same myeloma to the human kappa light chain exon. These genes were transfected into mouse myeloma cell lines and transformed cells producing chimeric mouse-human antiphosphocholine antibodies were thus developed.

[0009] Morrison, S. L. et al., European Patent Publication No. 173494 (published March 5, 1986), disclose chimeric "receptors" (e.g. antibodies) having variable regions derived from one species and constant regions derived from another. Mention is made of utilizing cDNA cloning to construct the genes, although no details of cDNA cloning or priming are shown. (see pp 5, 7 and 8).

[0010] Boulianne, G. L. et al., Nature, 312: 643 (December 13, 1984), also produced antibodies consisting of mouse variable regions joined to human constant regions. They constructed immunoglobulin genes in which the DNA segments encoding mouse variable regions specific for the hapten trinitrophenyl (TNP) were joined to segments encoding human μ and κ constant regions. These chimeric genes were expressed as functional TNP binding chimeric IgM.

[0011] For a commentary on the work of Boulianne et al. and Morrison et al., see Munro, Nature, 312: 597 (December 13, 1984), Dickson, Genetic Engineering News, 5, No. 3 (March 1985), or Marx, Science, 229: 455 (August 1985).

[0012] Neuberger, M. S. et al., Nature, 314: 268 (March 25, 1986), also constructed a chimeric heavy chain immunoglobulin gene in which a DNA segment encoding a mouse variable region specific for the hapten 4-hydroxy-3-nitrophenylacetate (NP) was joined to a segment encoding the human ϵ region. When this chimeric gene was transfected into the J558L cell line, an antibody was produced which bound to the NP hapten and had human IgE properties.

[0013] Neuberger, M. S. et al., have also published work showing the preparation of cell lines that secrete hapten-specific antibodies in which the Fc portion has been replaced either with an active enzyme moiety (Williams, G. and Neuberger, M. S. Gene 43:319, 1986) or with a polypeptide displaying c-myc antigenic determinants (Nature, 312:604, 1984).

- [0014] Nauberger, M. *et al.*, PCT publication WO 86/01533, (published March 13, 1986) also disclose production of chimeric antibodies (see p. 5) and suggests, among the technique's many uses the concept of "class switching" (see p. 6).
- [0015] Taniguchi, M., in European Patent publication No. 171 496 (published February 19, 1985) discloses the production of chimeric antibodies having variable regions with tumor specificity derived from experimental animals, and constant regions derived from human. The corresponding heavy and light chain genes are produced in the genomic form, and expressed in mammalian cells.
- [0016] Takeda, S. *et al.*, *Nature*, **314**: 452 (April 4, 1985) have described a potential method for the construction of chimeric immunoglobulin genes which have intron sequences removed by the use of a retrovirus vector. However, an unexpected splice donor site caused the deletion of the V region leader sequence. Thus, this approach did not yield complete chimeric antibody molecules.
- [0017] Cabilly, S. *et al.*, *Proc. Natl. Acad. Sci., USA*, **81**: 3273-3277 (June 1984), describe plasmids that direct the synthesis in *E. coli* of heavy chains and/or light chains of anti-carcinoembryonic antigen (CEA) antibody. Another plasmid was constructed for expression of a truncated form of heavy chain (Fd) fragment in *E. coli*. Functional CEA-binding activity was obtained by *in vitro* reconstitution, in *E. coli* extracts, of a portion of the heavy chain with light chain.
- [0018] Cabilly, S., *et al.*, European Patent Publication 125023 (published November 14, 1984) describes chimeric immunoglobulin genes and their presumptive products as well as other modified forms. On pages 21, 28 and 33 it discusses cDNA cloning and priming.
- [0019] Boss, M. A., European Patent Application 120694 (published October 3, 1984) shows expression in *E. coli* of non-chimeric immunoglobulin chains with 4-nitrophenyl specificity. There is a broad description of chimeric antibodies but no details (see p. 9).
- [0020] Wood, C. R. *et al.*, *Nature*, **314**: 446 (April, 1985) describe plasmids that direct the synthesis of mouse anti-NP antibody proteins in yeast. Heavy chain μ antibody proteins appeared to be glycosylated in the yeast cells. When both heavy and light chains were synthesized in the same cell, some of the protein was assembled into functional antibody molecules, as detected by anti-NP binding activity in soluble protein prepared from yeast cells.
- [0021] Alexander, A. *et al.*, *Proc. Natl. Acad. Sci., USA*, **79**: 3260-3264 (1982), describe the preparation of a cDNA sequence coding for an abnormally short human Ig gamma heavy chain (OMM gamma³ HCD serum protein) containing a 19- amino acid leader followed by the first 15 residues of the V region. An extensive internal deletion removes the remainder of the V and the entire C_H1 domain. This is cDNA coding for an internally deleted molecule.
- [0022] Dolby, T. W. *et al.*, *Proc. Natl. Acad. Sci., USA*, **77**: 6027-6031 (1980), describe the preparation of a cDNA sequence and recombinant plasmids containing the same coding for μ and κ human immunoglobulin polypeptides. One of the recombinant DNA molecules contained codons for part of the C_H3 constant region domain and the entire 3' noncoding sequence.
- [0023] Seno, M. *et al.*, *Nucleic Acids Research*, **11**: 719-726 (1983), describe the preparation of a cDNA sequence and recombinant plasmids containing the same coding for part of the variable region and all of the constant region of the human IgE heavy chain (epsilon chain).
- [0024] Kurokawa, T. *et al.*, *ibid.*, **11**: 3077-3085 (1983), show the construction, using cDNA, of three expression plasmids coding for the constant portion of the human IgE heavy chain.
- [0025] Liu, F. T. *et al.*, *Proc. Natl. Acad. Sci., USA*, **81**: 5369-5373 (September 1984), describe the preparation of a cDNA sequence and recombinant plasmids containing the same encoding about two-thirds of the C_H2, and all of the C_H3 and C_H4 domains of human IgE heavy chain.
- [0026] Tsujimoto, Y. *et al.*, *Nucleic Acids Res.*, **12**: 8407-8414 (November 1984), describe the preparation of a human V lambda cDNA sequence from an Ig lambda-producing human Burkitt lymphoma cell line, by taking advantage of a cloned constant region gene as a primer for cDNA synthesis.
- [0027] Murphy, J., PCT Publication WO 83/03971 (published November 24, 1983) discloses hybrid proteins made of fragments comprising a toxin and a cell-specific ligand (which is suggested as possibly being an antibody).
- [0028] Tan, *et al.*, *J. Immunol.*, **135**: 8564 (November, 1985), obtained expression of a chimeric human-mouse immunoglobulin genomic gene after transfection into mouse myeloma cells.
- [0029] Jones, P. T. *et al.*, *Nature*, **321**: 552 (May 1986) constructed and expressed a genomic construct where CDR domains of variable regions from a mouse monoclonal antibody were used to substitute for the corresponding domains in a human antibody.
- [0030] Sun, L. K. *et al.*, *Hybridoma* **5** suppl. 1: 517 (1986), describes a chimeric human/mouse antibody with potential tumor specificity. The chimeric heavy and light chain genes are genomic constructs and expressed in mammalian cells.
- [0031] Sahagan *et al.*, *J. Immunol.*, **137**: 1066-1074 (August 1986) describe a chimeric antibody with specificity to a human tumor associated antigen, the genes for which are assembled from genomic sequences.
- [0032] For a recent review of the field see also Morrison, S. L., *Science*, **229**: 1202-1207 (September 20, 1985) and Oi, V. T., *et al.*, *BioTechniques*, **4**: 214 (1986).
- [0033] The Oi, *et al.*, paper is relevant as it argues that the production of chimeric antibodies from cDNA constructs

in yeast and/or bacteria is not necessarily advantageous.

[0034] See also Commentary on page 835 in Biotechnology 4 (1986).

SUMMARY OF THE INVENTION

[0035] The invention provides a novel approach for producing genetically engineered antibodies of desired variable region specificity and constant region properties through gene cloning and expression of light and heavy chains. The cloned immunoglobulin gene products can be produced by expression in genetically engineered organisms.

[0036] The application of chemical gene syntheses, recombinant DNA cloning, and production of specific immunoglobulin chains in various organisms provides an effective solution for the efficient large scale production of human monoclonal antibodies. The invention also provides a solution to the problem of class switching antibody molecules, so as to readily prepare immunoglobulins of a certain binding specificity of any given class.

[0037] According to the present invention there are provided polynucleotide molecules comprising a prokaryotic promoter region in operable linkage to a dicistronic transcription unit, the unit encoding a heavy chain immunoglobulin, or a fragment thereof, and a light chain immunoglobulin, the heavy and light chains being separately operably linked to a sequence coding for a polypeptide secretion signal.

[0038] The invention also relates to such recombinant molecules in vehicles such as plasmid vectors, capable of expression in desired prokaryotic hosts.

[0039] The invention provides prokaryotic hosts capable of producing, by culture, chimeric antibodies and methods of using these hosts.

[0040] The invention may be used to provide chimeric immunoglobulin individual chains, whole assembled molecules, and immunoglobulin fragments (such as Fab) having human constant regions and non-human variable regions, wherein both variable regions have the same binding specificity.

[0041] Among other immunoglobulin chains and/or molecules that can be provided by the invention are:

(a) a complete functional, immunoglobulin molecule comprising:

- (i) two identical chimeric heavy chains comprising a non-human variable region and human constant region and
- (ii) two identical all (i.e. non-chimeric) human light chains.

(b) a complete, functional, immunoglobulin molecule comprising:

- (i) two identical chimeric heavy chains comprising a non-human variable region and a human constant region, and
- (ii) two identical all (i.e. non-chimeric) non-human light chains.

(c) a monovalent antibody, i.e., a complete, functional immunoglobulin molecule comprising:

- (i) two identical chimeric heavy chains comprising a non-human variable region and a human constant region, and
- (ii) two different light chains, only one of which has the same specificity as the variable region of the heavy chains. The resulting antibody molecule binds only to one end thereof and is therefore incapable of divalent binding;

(d) an antibody with two different specificities, i.e., a complete, functional immunoglobulin molecule comprising:

- (i) two different chimeric heavy chains, the first one of which comprises a non-human variable region and a human constant region and the second comprises a different non-human variable region, and a human constant region, and
- (ii) two different chimeric light chains, the first one of which comprises a non-human variable region having the same specificity as the first heavy chain variable region, and a human constant region, and the second comprises a non-human variable region having the same specificity as the second heavy chain variable region, and a human constant region.

[0042] The resulting antibody molecule binds to two different antigens.

[0043] The invention also provides for the production of functionally active chimeric immunoglobulin fragments secreted by prokaryotic or eukaryotic hosts or fully folded and reassembled chimeric immunoglobulin chains.

[0044] Genetic sequences, especially cDNA sequences, coding for the aforementioned combinations of chimeric

chains or of non-chimeric chains are also provided herein.

[0045] The use of cDNA sequences is particularly advantageous over genomic sequences (which contain introns), in that cDNA sequences can be expressed in bacteria or other hosts which lack RNA splicing systems.

[0046] Among preferred specific antibodies are those having specificities to cancer-related antigens.

BRIEF DESCRIPTION OF THE FIGURES

[0047] FIGURE 1 shows the DNA rearrangements and the expression of immunoglobulin μ and γ heavy chain genes. This is a schematic representation of the human heavy chain gene complex, not shown to scale. Heavy chain variable V region formation occurs through the joining of V_H , D and J_H gene segments. This generates an active μ gene. A different kind of DNA rearrangement called "class switching" relocates the joined V_H , D and J_H region from the μ constant C region to another heavy chain C region (switching to γ is diagrammed here). The scheme emphasizes that the J region is a common feature of all expressed heavy chain genes. The J region is also a common feature of expressed light chain genes.

[0048] FIGURE 2 shows the known nucleotide sequences of human and mouse J regions. Consensus sequences for the J regions are shown below the actual sequences. The oligonucleotide sequence below the mouse κ J region consensus sequence is a Universal Immunoglobulin Gene (UIG) oligonucleotide which is used in the present invention.

[0049] FIGURE 3 shows a scheme noting the use of the UIG oligonucleotide primer for the synthesis of cDNA complementary to the variable region of immunoglobulin messenger RNA, or the use of oligo-dT as a primer for cDNA synthesis, followed by *in vitro* mutagenesis.

[0050] FIGURE 4 shows the synthesis and analysis of human IgG1 genes, including three isolated clones (A, B), one of which (pGMH-6) is utilized as a cloning vector (B). A 1.5 kb deletion of pBR322 sequence between Bam HI and Pvu II is marked. Not to scale.

[0051] FIGURE 5 shows the cloning vector pQ23, a modified pBR322, useful for cDNA cloning at the KpnI site. This vector also contains the useful restriction enzyme sites Bgl II plus Sal I. Not to scale.

[0052] FIGURE 6 shows in A, the synthesis and analysis of human light chain κ genes. The Figure also shows in B, (not in scale) construction of a human C_K region cloning vector pING2001.

[0053] FIGURE 7 shows primers designed for immunoglobulin V region synthesis. (A) shows the heavy chain J-C regions and primers. A DNA version of each mouse J heavy region is shown directly above primers designed from that sequence. Mouse J regions are 5' to 3', left to right, while primers are 3' to 5', left to right. Primer names are included in brackets, and numbers of nucleotides (N) and number of mismatches with each J_H region are listed to the right. Primers which introduce a Bst EI site are underlined. (B) shows the light chain J regions and primers. The same as for (A) except for light chains. Primers designed to introduce a Bgl II site are underlined, as is the Bcl I site present in pING2016E. (C) shows mouse variable region consensus UIG primers. The actual primer sequence is shown below that consensus sequence. The human C_K Hind III vector pGML60 is shown below. (D) shows a mouse gamma 2a J/C junction primer.

[0054] FIGURE 8 shows the synthesis of heavy chain V region module genes using oligonucleotide primed cDNA synthesis. Not to scale.

[0055] FIGURE 9 shows the construction of hybrid mouse-human immunoglobulin genes. Panel A shows construction of a heavy chain gene. Stippled regions show C region modules, while hatched or black regions show V region modules. Not to scale.

[0056] FIGURE 10 shows the construction of cDNA cloning-expression shuttle vectors for mammalian cells. The vectors pING2003 and pING2003E are derived from pL1, pUC12, pSV2-neo and M8- α RX12. Stippled regions indicate mouse heavy chain enhancer DNA, hatched regions indicate SV-40 DNA from pL1, and cross-hatched regions indicate SV-40 DNA from pSV2-neo. In the vectors pING2003 and pING2003E, thick lines represent pBR322 DNA from pSV2-neo, while thin lines represent pUC12 DNA. Arrows indicate the locations and directions of SV-40 early region promoters, and indicates a complete SV-40 intron sequence. Not to scale.

[0057] FIGURE 11 shows the construction of the heavy chain expression plasmid pING2006E. Arrows show SV-40 promoter locations and directions of transcription. Hatched and black areas show mouse V region modules, while stippled areas show human C region modules. Not to scale.

[0058] FIGURE 12 shows the structure of the chimeric anti-hepatitis heavy chain genes in the expression plasmids pING2006E and pING2012E. Panel A shows the structure of mouse-human chimeric anti-hepatitis heavy chain genes. The structure of human IgG1 mRNA and cDNA is shown in A.a. The human heavy chain constant region cDNA clone pGMH-6 and the mouse heavy chain variable region cDNA clones pBS13-1 and pJ3-11 were used to make the hybrid gene used in pING2006E. Hatched gene blocks indicate mouse variable region sequences, while open gene blocks show human IgG1 constant region sequences. Panel B shows the nucleotide sequence of the anti-hepatitis B heavy chain variable region in pING2006E and pING2012E. pING2012E was constructed by first inserting a Bgl II site at the

Sall site of pING1202 (See Figure 16) to form pING1202BgIII. The chimeric heavy chain gene from this plasmid was inserted into the expression vector pING2003E, resulting in pING2012E. pING2012E differs from pING 2006E in the region immediately upstream of the initiator ATG. Underlined nucleotides denote human J region sequences from the cDNA clone pGMH-6. Asterisk amino acid 117 indicates a single change at this site from mouse to human sequence (Ala to Ser) introduced in the chimeric gene J region. Sequencing was by the Sanger method on plasmid (open circle) and M13 (closed circle) templates.

[0059] FIGURE 13 shows in panel A the J-C junction region nucleotide sequence in light chain clones derived from pING2001 (pMAK-3, pING2013E, pING2007E, pING2010E-gpt and pING2014E-gpt). The J region sequence originating from pk2-3 is marked human JK4. The G nucleotide not predicted by genomic sequencing is marked with an asterisk. The oligonucleotide primer (K2-4BCL1) used to modify this sequence is shown below the human JK4 sequence. Panel B diagrams the method of site-directed mutagenesis used to make pING2016E-gpt. Not to scale.

[0060] FIGURE 14 Gene copy number of the transfected sequences in two transformants. nDNA from 2AE9, 2BH10 were digested with the enzymes indicated. The concentration of DNA is titrated down across the lanes with the amount indicated above them. The probe contains human C gamma 1 sequences (pmvHc24 ApaI-BamHI). The reference is germ-line or GM2146 nDNA digested with ApaI. The 3' ApaI site is 2 bp beyond the site of poly(A) addition (3).

[0061] FIGURE 15 shows the nucleotide sequence of the V region of the L6 V_H cDNA clone pH3-6a. The sequence was determined by the dideoxytermination method using M13 subclones of gene fragments (shown below). Open circles denote amino acid residues confirmed by peptide sequence. A sequence homologous to D_{5P2} in the CDR3 region is underlined.

[0062] FIGURE 16 shows the nucleotide sequence of the V region of the L6 V_K cDNA clone pL3-12a. The oligonucleotide primer used for site-directed mutagenesis is shown below the J_{K5} segment. Open circles denote amino acid residues confirmed by peptide sequence.

[0063] FIGURE 17 shows the construction of chimeric L6-V_H plus human C gamma expression plasmids. Panel (a) shows the sequences of the BAL-31 deletion clones M13mp19-C1-delta 4 (C1-delta 4) and M13mp19-C1-delta 21(C1-delta 21). The 5' end of the cDNA clone, pH3-6a, is denoted with an arrow. M13 sequences are underlined. The oligonucleotide primer used for this experiment is H3-6a (5'- GACTGCACCACTGG-3'), which primes in FR1 near the mature N terminus. Panel (b) shows the strategy for site-directed mutagenesis of 1 ug of clones C1-delta 4 and C1-delta 21, each annealed to 20 ng of the 31-mer oligonucleotide MJH2-ApaI. Complementary strand synthesis with the Klenow fragment of DNA polymerase was at room temperature for 30 min, then 15°C for 72 hours. Transfected phage plaques were adsorbed to nitrocellulose, fixed with NaOH, and hybridized to ³²P-labelled MJH2-ApaI oligonucleotide at 65°C, 18 hours, in 4xTBS (0.6 M NaCl, 0.04 M Tris-HCl (pH 7.4), 0.004 M EDTA) plus 10% dextran sulfate. Final wash of the filters was at 65°C, 4xSSPE, 0.1% SDS for 15 min. (Maniatis, T., et al., Molecular Cloning: A Laboratory Manual, 1982). Positive plaques were detected by overnight exposure to Kodak XAR film, and were directly picked for growth and restriction enzyme analysis of RF DNA. Mismatches of the MJH2-ApaI oligonucleotide to the mouse C_{H1} are denoted, resulting in the coding changes shown below the oligonucleotide. Panel (c) shows the strategy of the substitution of each of the mutagenized L6-V_H modules for the resident V_H of the chimeric expression plasmid pING2012 to generate pING2111 and pING2112.

[0064] FIGURE 18 shows the construction of the chimeric L6 expression plasmid pING2119. The Sall to BamHI fragment from pING2100 is identical to the Sall to BamHI A fragment from pING2012E.

[0065] FIGURE 19 shows the modification of the V_K gene and its use in constructing light chain and heavy plus light chain expression plasmids.

(a) Deletion of the oligo d[GC] segment 5' of V_K of L6. The oligonucleotide is a 22-mer and contains a Sall site. The 3 mismatches are shown. The V_K gene, after mutagenesis, is joined as a Sall-HindIII fragment to the human C_K module. The expression plasmid thus formed is pING2119.

(b) pING2114, a heavy plus light chain expression plasmid. The expression plasmid pING2114 contains the L6 heavy chain chimeric gene from pING2111 and the chimeric light chain from pING2119 (bold line).

[0066] FIGURE 20 shows a summary of the sequence alterations made in the construction of the L6 chimeric antibody expression plasmids. Residues underlined in the 5' untranslated region are derived from the cloned mouse kappa and heavy-chain genes. Residues circled in the V/C boundary result from mutagenesis operations to engineer restriction enzyme sites in this region. Residues denoted by small circles above them in the L6 heavy-chain chimera also result from mutagenesis. They are silent changes.

[0067] FIGURE 21 shows the 2H7 V_H sequence. The V_H gene contains J_{H1} sequences and DSP.2 sequence elements. Small circles above the amino acid residues are those that matched to peptide sequences.

[0068] FIGURE 22 shows the 2H7 V_L sequence. The V_K gene contains J_{K5} sequences. A 22-mer oligonucleotide was used to place a Sall site 5' of the ATG initiator codon. Small circles above the amino acid residues are those that matched to peptide sequences.

[0069] FIGURE 23 shows the chimeric immunoglobulin gene expression plasmids of the 2H7 specificity. One gene plasmids are pING2101 (V_H ,neo), pING2105 (V_K ,neo) and pING2107 (V_K ,gpt). The others are two-gene plasmids. Their construction involved the ligation of the larger *NdeI* fragments of pING2101 and pING2107 to linearized pING2106 partially digested with *NdeI*. pHL2-11 and pHL2-26 were obtained from pING2101 and pING2105; pLL2-25 was obtained from pING2107 and pING2106.

[0070] FIGURE 24 shows a summary of the nucleotide changes introduced in the V_H and V_K in the construction of the chimeric plasmids. The cognate V_H and V_K nucleotide residues in the 5' and are underlined. Circles residues in the J-C junctions are derived from the human C modules.

[0071] FIGURE 25 shows the strategy used to fuse the mature L6 chimeric light chain sequence to the yeast invertase signal sequence and shortened PGK promoter. The open double line represents yeast invertase signal sequence DNA. The solid double line represents yeast PGK DNA, -> represents the PGK promoter, -I represents the PGK terminator, RF = Replicative Form. pING1225 was derived by fusing human C_κ DNA to the PGK promoter. pING1149 was derived by fusing the yeast invertase signal sequence to the yeast PGK promoter. (A) shows the strategy for introduction by *in vitro* mutagenesis of an *AatII* site in the signal sequence processing site. (B) shows the DNA sequence of the single-stranded mutagenesis primer and the corresponding unmutagenized DNA sequence. (C) shows the strategy used to construct a plasmid containing the mature light chain sequence fused to the invertase signal sequence and shortened PGK promoter.

[0072] FIGURE 26 shows the strategy used to fuse the mature L6 chimeric heavy chain sequence to the yeast invertase signal sequence and shortened PGK promoter. pING1288 contains the chimeric heavy chain gene with the variable region from the 2H7 mouse monoclonal antibody (see example IV). All symbols are as defined in legend for Figure 25. (A) shows the strategy for introduction by *in vitro* mutagenesis of an *SstI* site in the signal sequence processing site. (B) shows the DNA sequence of the single-stranded mutagenesis primer and the corresponding unmutagenized DNA sequence. (C) shows the strategy used to construct a plasmid containing the mature heavy chain sequence fused to the invertase signal sequence and shortened PGK promoter.

[0073] FIGURE 27 shows the strategy used to remove non-yeast 3' untranslated DNA sequences from the L6 chimeric light chain gene and to construct a plasmid containing the light chain gene fused to the invertase signal sequence and shortened PGK promoter in which all sequences are either known by DNA sequence analysis or proven to be functional. pBR322NA is derived from pBR322 by deletion of DNA from *NdeI* to *AuaI*. Symbols are as defined in legend for Figure 25.

[0074] FIGURE 28 shows the strategy used to remove non-yeast 3' untranslated DNA sequence from the L6 chimeric heavy chain gene and to construct a plasmid containing the heavy chain gene fused to the invertase signal sequence and shortened PGK promoter in which all sequences are either known by DNA sequence analysis or proven to be functional. Symbols are as defined in legend for Figure 25.

[0075] FIGURE 29 shows the strategy used to clone the L6 chimeric light chain gene fused to the invertase signal sequence and shortened PGK promoter into yeast-*E. coli* shuttle vectors containing the PGK transcription termination-polyadenylation signal, yeast replication sequences, and genes for selection of transformants. Symbols are as defined in legend for Figure 25.

[0076] FIGURE 30 shows the strategy used to clone the L6 chimeric heavy chain gene fused to the invertase signal sequence and shortened PGK promoter into yeast-*E. coli* shuttle vectors containing the PGK transcription termination-polyadenylation signal, yeast replication sequences, and genes for selection of transformants. Symbols are as defined in legend for Figure 25.

[0077] FIGURE 31 shows a schematic diagram of the structure of human IgG1.

[0078] FIGURE 32(A) shows the strategy used to introduce a stop codon and *BclI* site into the hinge region of human gamma 1. (B) shows the DNA sequence of the single-stranded primer used for *in vitro* mutagenesis of the gamma-1 hinge region and the corresponding unmutagenized sequence. Vertical arrows represent inter-chain disulfide bonds. Symbols are as defined in legend for Figure 25.

[0079] FIGURE 33 shows the strategy used to fuse the L6 chimeric heavy chain gene containing a stop codon in the hinge region (Fd chain) to the yeast invertase signal sequence and shortened PGK promoter. Symbols are as defined in legend for Figure 25.

[0080] FIGURE 34 shows the strategy used to remove non-yeast 3' untranslated sequences from the L6 chimeric Fd chain and to construct a plasmid containing the Fd chain fused to the invertase signal sequence and shortened PGK promoter in which all sequences are either known by DNA sequence analysis or proven to be functional. Symbols are as defined in legend for Figure 25.

[0081] FIGURE 35 shows the strategy used to clone the L6 chimeric Fd chain gene fused to the invertase signal sequence and shortened PGK promoter into yeast-*E. coli* shuttle vectors containing the PGK transcription termination-polyadenylation signal, yeast replication sequence, and genes for selection of transformants. Symbols are as defined in legend for Figure 25.

[0082] FIGURE 36(A) shows the nucleotide sequence surrounding the N-terminus of the *Erwinia caratovora* pB

gene (Lei, S.P., *et al.*, *J. Bacteriol.*, (1987, in press)). The *NdeI* and *HaeIII* sites used in cloning are shown. The arrow indicates the leader peptidase cleavage site for peptate lyase. (B) shows the cloning strategy for construction of the *pelB* leader cartridge. pSS1004 contains a 1.9 kb *DraI* fragment cloned into the *SmaI* site of pUC8. Symbols are defined in the legend for Figure 39.

[0083] FIGURE 37 shows the construction of light chain expression plasmids pRR177-8, pRR180, pRR190, and pRR191. In addition to the plasmids described in the text, M13mp18 and pIT181 were used. pIT181 contains the mature light chain gene fused directly following the ATG initiation codon of the *araB* gene in pIT2 (see Figure 40).

[0084] FIGURE 38 shows the construction of Fd expression plasmids pRR178-5, pRR186, and pRR196.

[0085] FIGURE 39 shows the restriction maps of the light chain and Fd gene cassette in pFK100, pFK101, pFK102, pFK103, and pFK104. These plasmids were constructed as described in the text using plasmids outlined in Figure 37 and 38. The arrow indicates the direction of transcription from the *lac* promoter. *E. caratovora* and eukaryotic non-coding sequences are shown as open bars. The *pelB* leader sequence is cross-hatched and the closed bar represents the antibody genes Fd and light chain (K).

[0086] FIGURE 40(A) shows the construction of a vector for arabinose inducible Fab expression. Plasmid pIT2 (Mason and Ray, *Nucl. Acids Res.*, 14:5693 (1986)) is a 6431 bp plasmid encoding the *araC* gene, the *araB* promoter, and a portion of the *araB* gene from pING1 (Johnston, S., *et al.*, *Gene* 34:134 (1985)) in a derivative of pBR322. An *NcoI* site has been engineered at the ATG initiation codon of the *araB* gene. (B) shows the introduction of the *laci* gene into pFK102.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

INTRODUCTION

[0087] Generally, antibodies are composed of two light and two heavy chain molecules. Light and heavy chains are divided into domains of structural and functional homology. The variable regions of both light (V_L) and heavy (V_H) chains determine recognition and specificity. The constant region domains of light (C_L) and heavy (C_H) chains confer important biological properties such as antibody chain association, secretion, transplacental mobility, complement binding, and the like.

[0088] A complex series of events leads to immunoglobulin gene expression in B cells. The V region gene sequences conferring antigen specificity and binding are located in separate germ line gene segments called V_H , D and J_H ; or V_L and J_L . These gene segments are joined by DNA rearrangements to form the complete V regions expressed in heavy and light chains respectively (Figure 1). The rearranged, joined (V_L - J_L and V_H -D- J_H)V segments then encode the complete variable regions or antigen binding domains of light and heavy chains, respectively.

DEFINITIONS

[0089] Certain terms and phrases are used throughout the specification and claims. The following definitions are provided for purposes of clarity and consistency.

1. Expression vector - a plasmid DNA containing necessary regulatory signals for the synthesis of mRNA derived from gene sequences, which can be inserted into the vector.
2. Module vector - a plasmid DNA containing a constant or variable region gene module.
3. Expression plasmid - an expression vector that contains an inserted gene, such as a chimeric immunoglobulin gene.
4. Gene cloning - synthesis of a gene, insertion into DNA vectors, and identification by hybridization and the like.
5. Transfection - the transfer of DNA into mammalian cells.
6. Promoter region - a nucleotide sequence which provides a cell with the regulatory sequences needed to express an operably linked cistron or operon.
7. Secretion signal - a polypeptide present at the N-terminus of a chimeric immunoglobulin chain useful in aiding in the secretion of the chain to the outside of the host. Also called "leading peptide," or "leader."

GENETIC PROCESSES AND PRODUCTS

[0090] The invention provides a novel approach for the cloning and production of human antibodies with desired specificity. Generally, the method combines five elements:

- (1) Isolation of messenger RNA (mRNA) from B cell hybridoma lines producing monoclonal antibodies against specific antigens, cloning and cDNA production therefrom;

(2) Preparation of Universal Immunoglobulin Gene (UIG) oligonucleotides, useful as primers and/or probes for cloning of the variable region gene segments in the light and heavy chain mRNA from specific human or non-human hybridoma cell lines, and cDNA production therefrom;

(3) Preparation of constant region gene segment modules by cDNA preparation and cloning, or genomic gene preparation and cloning;

(4) Construction of complete heavy or light chain coding sequences by linkage of the cloned specific immunoglobulin variable region gene segments of part (2) above to cloned human constant region gene segment modules;

(5) Expression and production of light and heavy chains in selected prokaryotic hosts, either in separate fermentations followed by assembly of antibody molecules in vitro, or through production of both chains in the same cell.

[0091] The invention employs cloned hybridoma B cell lines producing monoclonal antibodies of defined specificity for the isolation of mRNA for cDNA cloning. Because many lymphoid cell lines contain highly active nucleases which degrade mRNA during isolation, the invention uses mRNA preparation methods specifically developed for the isolation of intact mRNA from cells and tissues containing active nucleases. One such method yields total RNA preparations by intact mRNA disruption is an ethanol/perchlorate dry ice mixture which reduces nuclease action (Lizardi, P. M. et al., Anal. Biochem., **98**: 116 (1979)). This method gives intact translatable mRNA.

[0092] Other methods that have been used for this invention include extraction of cells with lithium chloride plus urea (Aufray, C., and Rougeon, F., Eur. J. Biochem., **107**: 303 (1980)) or guanidine thiocyanate (Chirgwin, J. M. et al., Biochemistry, **18**: 5294 (1979)) to prepare total RNA.

[0093] One universal feature of all expressed immunoglobulin light and heavy chain genes and messenger RNAs is the so-called J region (i.e. joining region, see Figure 1). Heavy and light chain J regions have different sequences, but a high degree of sequence homology exists (greater than 80%) within the heavy J_H regions or the J_{κ} light chain J regions. The invention provides consensus sequences of light and heavy chain J regions useful in the design of oligonucleotides (designated herein as UIGs) for use as primers or probes for cloning immunoglobulin light or heavy chain mRNAs or genes (Figures 2 or 7). Depending on the nature of design of a particular UIG, it may be capable of hybridizing to all immunoglobulin mRNAs or genes containing a single specific J sequence, such as UIG-MJH3 which detects only mouse J_{κ} sequences (Figure 7).

[0094] Another utility of a particular UIG probe may be hybridization to light chain or heavy chain mRNAs of a specific constant region, such as UIG-MJK which detects all mouse J_{κ} containing sequences (Figure 7). UIG design can also include a sequence to introduce a restriction enzyme site into the cDNA copy of an immunoglobulin gene (see Figure 7). The invention may, for example, utilize chemical gene synthesis to generate the UIG probes for the cloning of V regions in immunoglobulin mRNA from hybridoma cells making monoclonal antibodies of desired antigen specificities.

[0095] A multi-stage procedure is utilized for generating complete V + C region cDNA clones from hybridoma cell light and heavy chain mRNAs. In the first stage, the invention utilizes UIG probes as "primers" for reverse transcriptase copying of the complete V region and leader coding sequences of heavy and light chain mRNAs (Figure 3). The complementary strand of the primer extended cDNA is then synthesized, and this double-stranded cDNA is cloned in appropriate cDNA cloning vectors such as pBR322 (Gubler and Hoffman, Gene, **25**: 263 (1983)) or pQ23 (Figure 5; Maniatis, T. et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Publications, New York, page 224 (1982)). Clones are screened for specific hybridization with UIG oligonucleotide probes. Positive heavy and light chain clones identified by this screening procedure are mapped and sequenced to select those containing V region and leader coding sequences.

[0096] An alternative method is to make cDNA clones using oligo-dT as a primer, followed by selection of light and heavy chain clones by standard hybridization methods.

[0097] A second stage utilizes cloning of C region gene segments to form heavy and light chain module vectors. In one method cDNA clones of human heavy and light chain immunoglobulin mRNA are prepared. These cDNA clones are then converted into C region module vectors by site-directed mutagenesis to place a restriction site at a desired location near a boundary of the constant region. An alternative method utilizes genomic C region clones as the source for C region module vectors.

[0098] A third stage of cDNA cloning involves the generation of complete light and heavy chain coding sequences with linked V and C regions. The cloned V region segments generated as above are excised and ligated to light or heavy chain C region module vectors. For example, one can clone the complete human J_{κ} light chain C region and the complete human gamma C region. In addition, one can modify a human gamma 1 region and introduce a termination codon, thereby obtain a gene sequence which encodes the heavy chain portion of an Fab molecule.

[0099] The coding sequences having operationally linked V and C regions are then transferred into appropriate expression systems for expression in appropriate hosts, prokaryotic or eukaryotic. Operationally linked means in-frame joining of coding sequences to derive a continuously translatable gene sequence without alterations or interruptions of the triplet reading frame.

[0100] One particular advantage of using cDNA genetic sequences in the present invention is the fact that they code

continuously for immunoglobulin chains, either heavy or light. By "continuously" is meant that the sequences do not contain introns (i.e. are not genomic sequences, but rather, since derived from mRNA by reverse transcription, are sequences of contiguous exons). This characteristic of the cDNA sequences provided by the invention allows them to be expressible in prokaryotic hosts, such as bacteria, or in lower eukaryotic hosts, such as yeast.

[0101] Another advantage of cDNA cloning methods is the ease and simplicity of obtaining V region gene modules.

[0102] The term "non-human" as used in the invention is meant to include any animal other than a human, wherein an immune response can be generated which then leads to usable cells resulting in corresponding hybridomas or B cell clones obtained by viral transformation and the like. Such animals commonly include rodents such as the mouse or the rat. Because of ease of preparation and great availability, the mouse is at present the preferred, non-human animal. Mouse-mouse hybridomas are thus utilized as the preferred sources for heavy and light chain variable regions.

[0103] Preferably, the invention provides entire V and/or C region cDNA sequences. This means that the sequences code for substantially operable V and/or C regions, without lacking any major structural portions thereof.

[0104] The terms "constant" and "variable" are used functionally to denote those regions of the immunoglobulin chain, either heavy or light chain, which code for properties and features possessed by the variable and constant regions in natural non-chimeric antibodies. As noted, it is not necessary for the complete coding region for variable or constant regions to be present, as long as a functionally operating region is present and available.

[0105] A wide range of source hybridomas are available for the preparation of mRNA. For example, see the catalogue ATCC CELL LINES AND HYBRIDOMAS, December, 1984, American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland 20852, U.S.A., pages 5-9 and the ECACC Catalogue, 2nd Edition; PHLS CAMR Porton Down, Salisbury, Wilt; SP40JG, U.K. pages 30-35 and 40-46. Hybridomas secreting monoclonal antibodies reactive to a wide variety of antigens are listed therein, are available from the collection, and usable in the invention. Of particular interest are hybridomas secreting antibodies which are reactive with viral antigens, including Dengue complex specific (ATCC HB 114), Dengue type 1 virus (ATCC HB 47), Dengue type 2 virus (ATCC HB 46), Dengue type 3 virus (ATCC HB 49), Dengue type 4 virus (ATCC HB 48), Epstein-Barr receptor (ATCC HB 135), Flavivirus group (ATCC HB 112), hepatitis B surface antigen (ATCC CRL 8017 and 8018), herpes simplex type I (ATCC HB 8068), herpes simplex type II (ATCC HB 8067), influenza virus (ATCC CL 189), Influenza A virus, matrix protein (ATCC HB 84), Influenza A virus, nucleoprotein (ATCC HB 85), Influenza A Bangkok/179HA (ATCC HB 66), Influenza AWSN NP (ATCC HB 67), SV40 large T antigen (ATCC TIB 115), SV40 large T antigen, C-terminal end (ATCC TIB 117), and SV40 nonviral T antigen (ATCC TIB 116). Examples of other hybridomas include those secreting antibodies to tumor associated antigens or to human lymphocyte antigens, such as those reactive to human tumor-associated CEA, high mw (ATCC CRL 8019); human tumor-associated α -fetoprotein, IgG₁K - (ATCC HB 134); human B lymphocyte HLA-DR, monomorphic, IgG_{2a} (ATCC HB 104); human T lymphocyte T cell precursors, IgG₁ (ATCC CRL 8022); human T lymphocyte T cell subset, helper, IgG_{2a} (ATCC CRL 8002); T subset, suppressor/cytotoxic, human, IgG₁ (ATCC CRL 8013); T cell subset, suppressor/cytotoxic, human, IgG_{2a} (ATCC CRL 8014); T cells, peripheral, human, IgG₁ (ATCC CRL 8000); T cells, peripheral, human, IgG_{2a} (ATCC CRL 8001); thymocytes, "common," human, IgG₁ (ATCC CRL 8020).

[0106] These lines and others of similar nature can be utilized to copy the mRNA coding for variable region, using the U1G probes. Of particular interest are antibodies with specificity to human tumor antigens.

[0107] Expression vehicles include plasmids or other vectors. Preferred among these are vehicles carrying a functionally complete human constant heavy or light chain sequence having appropriate restriction sites engineered so that any variable heavy or light chain sequence with the appropriate cohesive ends can be easily inserted therinto. Human constant heavy or light chain sequence-containing vehicles are thus an important embodiment of the invention. These vehicles can be used as intermediates for the expression of any desired complete heavy or light chain in any appropriate host.

[0108] Among bacterial hosts which may be utilized as transformation hosts, *E. coli* K12 strain 294 (ATCC 31446) is particularly useful. Other microbial strains which may be used include *E. coli* X1776 (ATCC 31537). The aforementioned strains, as well as *E. coli* W3110 (ATCC 27325) and other enterobacteria such as *Salmonella typhimurium* or *Serratia marcescens*, and various *Pseudomonas* species may be used.

[0109] In general, plasmid vectors containing replicon and control sequences which are derived from species compatible with a host cell are used in connection with these hosts. The vector ordinarily carries a replication site, as well as specific genes which are capable of providing phenotypic selection in transformed cells. For example, *E. coli* is readily transformed using pBR322, a plasmid derived from an *E. coli* species (Bolivar, et al., *Gene*, 2: 95 (1977)). pBR322 contains genes for ampicillin and tetracycline resistance, and thus provides easy means for identifying transformed cells. The pBR322 plasmid or other microbial plasmids must also contain, or be modified to contain, promoters which can be used by the microbial organism for expression of its own proteins. Those promoters most commonly used in recombinant DNA construction include the beta-lactamase (penicillinase) and lactose (beta-galactosidase) promoter systems (Chang et al., *Nature*, 275: 615 (1978); Itakura et al., *Science*, 198: 1056 (1977)); and tryptophan promoter systems (Goeddel et al., *Nucleic Acids Research*, 8: 4057 (1980); EPO Publication No. 0036776). While these are the most commonly used, other microbial promoters have been discovered and utilized.

[0110] For example, a genetic construct for any heavy or light chimeric immunoglobulin chain can be placed under the control of the leftward promoter of bacteriophage lambda (P_L). This promoter is one of the strongest known promoters which can be controlled. Control is exerted by the lambda repressor, and adjacent restriction sites are known.

[0111] The expression of the immunoglobulin chain sequence can also be placed under control of other regulatory sequences which may be "homologous" to the organism in its untransformed state. For example, lactose dependent *E. coli* chromosomal DNA comprises a lactose or lac operon which mediates lactose digestion by elaborating the enzyme beta-galactosidase. The lac control elements may be obtained from bacteriophage lambda pLAC5, which is infective for *E. coli*. The lac promoter-operator system can be induced by IPTG.

[0112] Other promoter/operator systems or portions thereof can be employed as well. For example, arabinose, colicine E1, galactose, alkaline phosphatase, tryptophan, xylose, tac, and the like can be used. Other bacterial gene expression control elements can be utilized to achieve the expression of immunoglobulin proteins. For example, a gene with a bacterial secretion signal peptide coding region can be expressed in bacteria, resulting in secretion of the immunoglobulin peptide which was originally linked to the signal peptide.

[0113] Since an immunoglobulin cDNA is comprised only of sequences representing the mature mRNA encoding an antibody protein or its precursor, additional gene expression elements regulating transcription of the gene and processing of the RNA are required for optimal synthesis of immunoglobulin mRNA. These elements may include splice signals, as well as transcription promoters including inducible promoters enhancers, and termination signals. cDNA expression vectors incorporating such elements include those described by Okayama, H. and Berg, P., *Mol. Cell Biol.*, **3**: 280 (1983); Cepko, C. L. *et al.*, *Cell*, **37**: 1053 (1984); and Kaufman, R.J., *Proc. Natl. Acad. Sci., USA*, **82**: 669 (1985).

[0114] Different approaches can be followed to obtain complete H_2L_2 antibodies.

[0115] First, one can separately express the light and heavy chains followed by *in vitro* assembly of purified light and heavy chains into complete H_2L_2 IgG antibodies. The assembly pathways used for generation of complete H_2L_2 IgG molecules in cells have been extensively studied (see, for example, Scharff, M., *Harvey Lectures*, **68**: 125 (1974)). *In vitro* reaction parameters for the formation of IgG antibodies from reduced isolated light and heavy chains have been defined by Beychok, S., *Cells of Immunoglobulin Synthesis*, Academic Press, New York, page 69, 1979.

[0116] Second, it is possible to co-express light and heavy chains in the same cells to achieve intracellular association and linkage of heavy and light chains into complete H_2L_2 IgG antibodies. The coexpression can occur by using either the same or different plasmids in the same host.

[0117] Co-expression with the aid of secretion signals useful in bacteria, enable fully folded and assembled H_2L_2 immunoglobulins to be obtained.

[0118] Also, preparation of chimeric Fab fragments can be carried out by the methods of the invention.

[0119] The methods described herein can also be used to switch the class of any antibody of a given specificity and class to an antibody of the same specificity but of a different class, whether human or non-human. For example, human IgM antibodies can be transmutated to human IgG antibodies by preparing constructs containing human constant IgG cDNA or genomic sequences, linked to variable human cDNA sequences obtained from a cell producing the original IgM antibody. These constructs are then introduced into appropriate hosts and expressed.

POLYPEPTIDE PRODUCTS

[0120] The invention provides "chimeric" immunoglobulin chains, either heavy or light. A chimeric chain contains a constant region substantially similar to that present in the heavy chain of a natural human immunoglobulin, and a variable region having any desired antigenic specificity. The variable region is either from human or non-human origin.

[0121] The invention also provides immunoglobulin molecules having heavy and light chains associated so that the overall molecule exhibits desired binding and recognition properties. Various types of immunoglobulin molecules are provided: monovalent, divalent, dispecific (i.e., with different variable regions), molecules with chimeric heavy chains and non-chimeric light chains, or molecules with variable binding domains attached to peptide moieties carrying desired functions.

[0122] Antibodies having chimeric heavy chains of the same or different variable region binding specificity and non-chimeric (i.e., all human or all non-human) light chains, can be prepared by appropriate association of the needed polypeptide chains. These chains are individually prepared by the modular assembly methods of the invention.

[0123] Chimeric Fab fragments are also part of this invention.

USES

[0124] The antibodies that can be made by the invention having human constant region can be utilized for passive immunization, especially in humans, without negative immune reactions such as serum sickness or anaphylactic shock. The antibodies can, of course, also be utilized in prior art immunodiagnostic assays and kits, in labelled form for *in vitro* imaging, wherein the label can be a radioactive emitter, or an NMR contrasting agent such as a carbon-13 nucleus,

or an X-ray contrasting agent, such as a heavy metal nucleus. The antibodies can also be used *in vitro* localization of antigens by appropriate labelling.

[0125] The antibodies can be used for therapeutic purposes by themselves in complement mediated lysis or can be coupled to toxins or other therapeutic moieties.

[0126] Class switching of antibodies is useful when it is desired to change the association, aggregation or other properties of antibodies obtained from cell fusion or hybridoma technology. For example, most human-human monoclonals are of the IgM class, which are known for their ease of reduction and aggregation. Changing such antibodies to other antibody types, such as IgG, IgA, or IgE, is thus of great benefit.

[0127] Mixed antibody-enzyme molecules can be used for immunodiagnostic methods, such as ELISA. Mixed antibody-peptide effector conjugates can be used for targeted delivery of the effector moiety with a high degree of efficacy and specificity.

[0128] Having now generally described the invention, the same will be further understood by reference to certain specific examples which are included herein for purposes of illustration only and are not intended to be limiting unless otherwise specified.

EXPERIMENTAL

Materials and Methods

Tissue Culture Cell Lines

[0129] The human cell lines GM2146 and GM1500 were obtained from the Human Mutant Cell Repository (Camden, New Jersey) and cultured in RPMI1640 plus 10% fetal bovine serum (M. A. Bioproducts). The cell lines Sp2/0 and CRL 8017 were obtained from the American Type Culture Collection and grown in Dulbecco's Modified Eagle Medium (DMEM) plus 4.5 g/l glucose (M. A. Bioproducts) plus 10% fetal bovine serum (Hyclone, Sterile Systems, Logan, Utah). Media were supplemented with penicillin/streptomycin (Irvine Scientific, Irvine, California).

Recombinant Plasmid and Bacteriophage DNAs

[0130] The plasmids pBR322, pL1 and pUC12 were purchased from Pharmacia P-L Biochemicals (Milwaukee, Wisconsin). The plasmids pSV2-neo and pSV2-gpt were obtained from BRL (Gaithersburg, Maryland), and are available from the American Type Culture Collection (Rockville, Maryland). pHu-gamma-1 is a subclone of the 8.3 Kb *Hind*III to *Bam*HI fragment of the human IgG1 chromosomal gene. A separate isolation of the human IgG1 chromosomal gene is described by Ellison, J. W. *et al.*, *Nucl. Acids Res.*, 10: 4071 (1982). MBalphaRX12 contains the 0.7 Kb *Xba*I to *Eco*RI fragment containing the mouse heavy chain enhancer from the J-C Intron region of the M603 chromosomal gene (Davis, M. *et al.*, *Nature*, 263: 733) inserted into M13mp10. G-tailed pUC9 was purchased from Pharmacia P-L. DNA manipulations involving purification of plasmid DNA by buoyant density centrifugation, restriction endonuclease digestion, purification of DNA fragments by agarose gel electrophoresis, ligation and transformation of *E. coli* were as described by Maniatis, T. *et al.*, *Molecular Cloning: A Laboratory Manual*, (1982). Restriction endonucleases and other DNA/RNA modifying enzymes were purchased from Boehringer-Mannheim (Indianapolis, Indiana), BRL, New England Biolabs (Beverly, Massachusetts) and Pharmacia P-L.

Oligonucleotide Preparation

[0131] Oligonucleotides were either synthesized by the triester method of Ho *et al.* (*Nucl. Acids Res.*, 10: 1755 (1982)), or were purchased from ELEN, Los Angeles, California. Tritylated, deblocked oligonucleotides were purified on Sephadex-G50, followed by reverse-phase HPLC with a 0-25% gradient of acetonitrile in 10mM triethylamine-acetic acid, pH 7.2, on a C18 uBondapak column (Waters Associates). Detritylation was in 80% acetic acid for 30 min., followed by evaporation *trice*. Oligonucleotides were labeled with [gamma-³²P]ATP plus T4 polynucleotide kinase.

RNA Preparation and Analysis

[0132] Total cellular RNA was prepared from tissue culture cells by the method of Aufray, C. and Rougeon, F. (*Eur. J. Biochem.*, 107: 303 (1980)) or Chirgwin, J. M. *et al.* (*Biochemistry*, 18: 5294 (1979)). Preparation of poly(A)⁺ RNA, methyl-mercury agarose gel electrophoresis, and "Northern" transfer to nitrocellulose were as described by Maniatis, T. *et al.*, *supra*. Total cellular RNA or poly(A)⁺ RNA was directly bound to nitrocellulose by first treating the RNA with formaldehyde (White, B. A. and Bancroft, F. C., *J. Biol. Chem.*, 257: 8569 (1982)). Hybridization to filterbound RNA was with nick-translated DNA fragments using conditions described by Margulies, D. H. *et al.* (*Nature*, 295: 166 (1982)).

or with ^{32}P -labelled oligonucleotide using 4xSSC, 10X Denhardt's, 100 ug/ml salmon sperm DNA at 37°C overnight, followed by washing in 4xSSC at 37°C.

cDNA Preparation and Cloning

[0133] Oligo-dT primed cDNA libraries were prepared from poly(A)⁺ RNA from GM1500 and GM2146 cells by the methods of Land, H. *et al.* (Nucl. Acids Res., 9: 2251 (1981)) and Gubler, V. and Hoffman, B. J., *Gene*, 25: 263 (1983), respectively. The cDNA libraries were screened by *in situ* hybridization (Maniatis, T., *supra*) with ^{32}P -labelled oligonucleotides using the conditions shown above, or with nick-translated DNA fragments using the conditions of de Lange *et al.* (Cell, 34: 891 (1983)).

Oligonucleotide Primer Extension and Cloning

[0134] Poly(A)⁺ RNA (20 ug) was mixed with 1.2 ug primer in 40 ul of 64mM KCl. After denaturation at 90°C for 5 min. and then chilling in ice, 3 units Human Placental Ribonuclease Inhibitor (BRL) was added in 3 ul of 1M Tris-HCl, pH 8.3. The oligonucleotide was annealed to the RNA at 42°C for 15 minutes, then 12 ul of .05M DTT, .05M MgCl₂, and 1 mM each of dATP, dTTP, dCTP, and dGTP was added. 2 ul of α - ^{32}P -dATP (400 Ci/mmol, New England Nuclear) was added, followed by 3 ul of AMV reverse transcriptase (19 units/ul, Life Sciences).

[0135] After incubation at 42°C for 105 min., 2 ul 0.5 M EDTA and 50 ul 10mM Tris, 1mM EDTA, pH 7.6 were added. Unincorporated nucleotides were removed by Sephadex G-50 spun column chromatography, and the RNA-DNA hybrid was extracted with phenol, then with chloroform, and precipitated with ethanol. Second strand synthesis, homopolymer tailing with dGTP or dCTP, and insertion into homopolymer tailed vectors was as described by Gubler and Hoffman, *supra*.

Site-Directed Mutagenesis

[0136] Single stranded M13 subclone DNA (1 ug) was combined with 20 ng oligonucleotide primer in 12.5 ul of Hin buffer (7 mM Tris-HCl, pH 7.6, 7 mM MgCl₂, 50 mM NaCl). After heating to 95°C in a sealed tube, the primer was annealed to the template by slowly cooling from 70°C to 37°C for 90 minutes. 2 ul dNTPs (1 mM each), 1 ul ^{32}P -dATP (10 uCi), 1 ul DTT (0.1 M) and 0.4 ul Klenow DNA Pol (2u, Boehringer Mannheim) were added and chains extended at 37°C for 30 minutes. To this was added 1 ul (10 ng) M13 reverse primer (New England Biolabs), and the heating/annealing and chain extension steps were repeated. The reaction was stopped with 2 ul of 0.5M EDTA, pH 8, plus 80 ul of 10 mM Tris-HCl, pH 7.6, 1 mM EDTA. The products were phenol extracted and purified by Sephadex G-50 spun column chromatography and ethanol precipitated prior to restriction enzyme digestion and ligation to the appropriate vector.

Transfection of Myeloma Tissue Culture Cells

[0137] A variation of the method of Ochi, A. *et al.* (Nature, 302: 340 (1983)) was used for protoplast fusion. 50 ml of bacteria at A₆₀₀ of 0.7 were converted to protoplasts by the method of Sandri-Goldini, R. M. *et al.* (Mol. Cell. Biol., 1: 743 (1981)), then diluted with 20 ml DMEM plus 10% FBS (final volume is 25 ml). Sp2/O cells were harvested, pelleted at 2,200 x g, washed, repelleted and resuspended in DMEM at 2-5x10⁶/ml. Bacterial protoplasts (10 ml) were mixed with 10x10⁶ Sp2/O cells and pelleted by centrifugation at 4,000 x g at 22°C for 20 min. After pipetting off the supernatant, the pellet was suspended in the remaining drop of medium by flicking the tube. 2ml of 10% DMSO, 37% (w/v) PEG6000 (Kodak) in DMEM was added dropwise with mixing over 45 sec. After 15 sec., 2 ml of 42% PEG6000 in DMEM was added over 45 sec. Complete DMEM (45 ml) was slowly added with mixing. Cells were pelleted at 2500 x g, then washed and pelleted thrice.

[0138] The electroporation method of Potter, H. *et al.* (Proc. Natl. Acad. Sci., USA, 81: 7161 (1984)) was used. After transfection, cells were allowed to recover in complete DMEM for 48-72 hours, then were seeded at 10,000 to 50,000 cells per well in 96-well culture plates in the presence of selective medium. G418 (GIBCO) selection was at 0.8 mg/ml, mycophenolic acid (Calbiochem) was at 6 ug/ml plus 0.25 mg/ml xanthine, and HAT (Sigma) was at the standard concentration.

Assays for Immunoglobulin Synthesis and Secretion

[0139] Secreted immunoglobulin was measured directly from tissue culture cell supernatants. Cytoplasmic protein extract was prepared by vortexing 1x10⁶ cells in 160 ul of 1% NP40, 0.15 M NaCl, 10 mM Tris, 1 mM EDTA, pH 7.6 at 0°C, 15 minutes, followed by centrifugation at 10,000 x g to remove insoluble debris.

[0140] Double antibody sandwich ELISA (Voller, A. *et al.*, in *Manual of Clinical Immunology*, 2nd Ed., Eds. Rose, N. and Friedman, H., pp. 359-371, 1980) using affinity purified antisera was used to detect specific immunoglobulins. For detection of human IgG, the plate-bound antiserum is goat anti-human IgG (KPL, Gaithersburg, Maryland) at 1/1000 dilution, while the peroxidase-bound antiserum is goat anti-human IgG (KPL or Tago, Burlingame) at 1/4000 dilution. For detection of human immunoglobulins kappa, the plate-bound antiserum is goat anti-human kappa (Tago) at 1/500 dilution, while the peroxidase-bound antiserum is goat anti-human kappa (Cappel) at 1/1000 dilution.

[0141] Antibodies binding hepatitis B surface antigen were detected using a commercial (Abbott, AUSAB) assay

EXAMPLES

[0142] The following examples, of which Example VII exemplifies the invention, show cDNA cloning for the modular assembly of antibody genes as used for the preparation of chimeric antibodies each having a human constant region and a non-human variable region. These examples outline the step-by-step process of preparing the chimeric antibodies.

EXAMPLE I: Human Antibody Constant Region Gene Modules and cDNA Expression Vectors

(1) Preparation of cDNA Clones, and Vehicles Containing Same, for Heavy Chain Human Constant Region

[0143] The cell line GM2146 was used as the source in mRNA preparation and cDNA cloning. This cell line secretes IgG, (Simmons, J. G. *et al.*, *Scand. J. Immunol.*, **14**: 1-13, 1981). Tests of this cell line indicated that it secretes IgA as well as IgG.

[0144] The cell line was cloned, and results indicated that five of six subclones secreted IgG only, while one of six subclones secreted IgA only. Poly(A)⁺ RNA was prepared from the cell line and a cDNA library was prepared from the poly(A)⁺ RNA by the method of Gubler, U. and Hoffman, B. J., *Gene*, **25**: 263-269 (1983). An initial plating of the cDNA transformed into E. coli strains HB101 and RR1 yielded a total of 1500 colonies, which were screened by hybridization to a HindIII to BamHI fragment of a genomic clone of human IgG1 (pHu-gamma-1). Four positive clones were found. A fragment containing the CH3 coding region of one of these clones, pGMH-3 (Figure 4), was used to rescreen the original library plus a new transformation of approximately 5000 colonies. Two of the largest clones, pGMH-6 and pGMH-15, were analyzed by restriction enzyme digestion (Figure 4). Both clones contained the entire constant region of human IgG1, although it was discovered that pGMH-6 had deleted approximately 1500 base pairs of pBR322 DNA, apparently without affecting the IgG1 cDNA sequences.

[0145] Clone pGMH-6 provided the IgG1 constant region module in the construction of cloning vectors for heavy chain variable region cloning.

(2) Preparation of cDNA Clones, and Vehicles Containing Same, for Light Chain Human Constant Region

[0146] A human cell line (GM1500) producing IgG2K was selected for the initial cloning phase. Poly(A)⁺ RNA prepared from GM1500 is active in *in vitro* translation using rabbit reticulocyte extracts. A cDNA library was prepared from this RNA by the method of Land *et al.*, *Nucl. Acids Res.*, **9**: 2251-2266 (1981), utilizing KpnI digested and dG-tailed pQ23 as the cloning vector (Figure 5). This vector contains BglII, KpnI and SacI sites inserted between the BamHI and SacI sites of pBR322.

[0147] In order to identify the cDNA clones generated from GM1500 RNA which correspond to light chain mRNA, a DNA probe, UIG-HuK, was synthesized and purified. The UIG-HuK oligonucleotide has the sequence 5'-AGCCACAGTTCGTTT-3', and is designed to hybridize to all functional human kappa mRNA species at the J-C junction. This probe was used to prime cDNA synthesis on GM1500 RNA in the presence of dideoxynucleotides and reverse transcriptase. From 1.2 ug of total GM1500 poly(A)⁺ RNA was used in this experiment, the entire J sequence and some of the V region was read, demonstrating that (1) GM1500 RNA is intact, (2) the kappa probe is of the correct sequence, and (3) GM1500 light chain mRNA contains J₄ sequences.

[0148] cDNA clones positive for hybridization to the light chain probe were selected. Since the probe hybridizes to the J-C junction, the most important point was to determine if the clones had complete constant region sequence in addition to the J region.

[0149] Insert sizes for the two largest kappa cDNA clones were 0.6 and 0.9 kb; restriction enzyme mapping indicated that the entire constant region coding sequence was present in both clones (Figure 6). The human kappa cDNA clone pk2-3 was used to make the light chain constant region vector pING2001 by inserting the Sau3A fragment comprising the human kappa constant and J regions into the BclI site of pBR325 (Figure 6B).

[0150] A variant of the human kappa cDNA clone was made by placing a HindIII site in the J region. This was carried out by *in vitro* mutagenesis using a J₄HINDIII oligonucleotide primer (Figure 7c). The resultant plasmid is pGML60.

[0151] A vector, pING2003, was constructed for the transfer and expression of cDNA sequences in mammalian cells (Figure 10). This vector was constructed from pUC12 and two plasmids containing SV40 sequences. pL1 provides an SV40 early region promoter and an SV40 late region splice sequence. pSV2-neo sequences provide a selectable marker for mammalian cell transformation and SV40 polyadenylation signal sequences. pUC12 provides a multiple cloning site for cDNA insertion.

[0152] The pING2003 vector has several useful restriction sites for modifications. These include a *Hind*III site useful for the insertion of enhancer sequences, and a *Hind*III to *Xba*I fragment useful for the insertion of alternate promoter sequences. This vector is useful in the expression of cDNA genes in mammalian cells.

10 Addition of Enhancer Element to pING2003

[0153] Immunoglobulin enhancer elements have been shown to enhance transcription of genes in their vicinity in stably transformed mouse myeloma cells by several hundred fold (Gillies, S. D. *et al.*, *Cell*, **33**: 717, 1983; and Banerji, J. *et al.*, *Cell*, **33**: 729, 1983). To facilitate expression of the mouse-human immunoglobulin genes in mouse myeloma cells, the mouse immunoglobulin heavy chain enhancer element was added to the cDNA expression vector pING2003 (Figure 10). The mouse heavy chain enhancer region DNA was isolated from an M13 subclone of mouse heavy chain genomic DNA (M8- α -RX12, Deans, R. J., unpublished). DNA isolated from a *Sall* plus *Eco*RI digestion of this subclone was modified with *Hind*III linkers and inserted into the *Hind*III site of pING2003, resulting in the new cDNA expression vector pING2003E. This vector is useful in the efficient expression of cDNA genes in mammalian cells, particularly mouse myeloma or hybridoma cell lines.

EXAMPLE II: Human-Mouse Chimeric Anti-HBsAg Antibody Chain

(1) Preparation of cDNA Clones and Vehicles Containing Same, for Heavy Chain Mouse Anti-HBsAg Variable Region.

[0154] The cell line CRL8017 was obtained from the ATCC and subcloned. Subclones were grown and tested for mouse IgG anti-hepatitis B binding activity using a commercially available anti-HBsAg detection kit. Three positive subclones were found. Poly(A)⁺ RNA was prepared from one of these subclones, and was fractionated on a methyl-mercury agarose gel. The RNA contained intact light chain and heavy chain mRNA's as inferred from specific hybridization to *kappa* UIG-MJK primer, and to the mouse heavy chain UIG-MJH3 probe (see Figure 7). In addition, the UIG-MJK primer was used for specific priming of anti-HBsAg poly(A)⁺ RNA in a dideoxy sequencing reaction. Sufficient sequence was read to show that a major *kappa* RNA of the anti-HBsAg cell line contains the J_H2 sequence.

[0155] The conditions for variable region cDNA synthesis were optimized by using heavy and light chain UIG primers on anti-HBsAg poly(A)⁺ RNA. Dideoxy chain extension experiments demonstrated that the mouse UIG-MJK primer and UIG-JH3 primer correctly primed *kappa* and heavy chain RNAs. When the reverse transcription was carried out in the absence of dideoxynucleotides, the main product using the *kappa* UIG-MJK primer was a 410±20 nucleotide fragment, while the main product using the heavy chain UIG-JH3 primer was a 430±30 nucleotide fragment. These correspond to the expected lengths of the variable and 5' untranslated regions of *kappa* and heavy chain immunoglobulin mRNAs. The conditions for the optimal priming of poly(A)⁺ RNA from CRL8017 cells should work well for poly(A)⁺ RNA isolated from any cell line producing a monoclonal antibody.

[0156] After determining optimal conditions for priming hybridoma mRNA with oligonucleotide primers, two oligonucleotides were designed and used for heavy chain V region cDNA synthesis. These two oligonucleotides are UIG-MJHSTEII(13) and UIG-MJH3 (Figures 7 and 8). It should be noted that the primer sequence was designed to introduce a *Bst*EII recognition site (GGTGACC) in the clone so that it could be joined at this site to the human IgG1 constant module at the analogous position at the latter's J region. In this case, the primer had a single G to U mismatch with the mouse mRNA sequence that uses the J_H3 coding sequence. The UIG-MJHSTEII(13) primer was 13 bases long and the mismatched residue was flanked by 7 matches 5' and 5 matches 3' of it. This was the 13-mer *Bst*EII primer. To assess the priming efficiency of the 13-mer *Bst*EII oligonucleotide, a 21-mer primer specific for mouse J_H3 (UIG-MJH3) was used. This primer had a perfect match for the 17 nucleotides on its 3' end.

[0157] These two primers and the J_H3 coding sequences are shown in Figure 8. The first strand cDNA products made via the 13-mer *Bst*EII and the 21-mer J_H3 primers included bands of approximately 430 nucleotides, which represented the entire V_H region. Under the standard priming conditions used, the priming efficiency of the 13-mer *Bst*EII was much less than that of the 21-mer J_H3. Accordingly, a cDNA library was generated from the first strand synthesis from each of these primers, using the method of Gubler and Hoffman, *supra*.

[0158] First, the 21-mer J_H3 library was screened with the 21-mer J_H3 oligonucleotide. Filter hybridization was done at 30°, overnight, according to de Lange, T. *et al.*, *Cell*, **34**: 891-900 (1983). The filters were then washed at 51° in 6 x SSC, 0.1% SDS. Five colonies were selected. The largest had an insert of approximately 460 bp. More significantly, it contained three restriction sites predicted from the known J_H3 sequence, which are present upstream of the primer

sequence. This clone, pJ3-11, was sequenced using the J_κ3 primer by the chain-termination method (Wallace, R. B. *et al.*, *Gene*, 16: 21-26 (1981)). The sequence obtained has the remaining J_κ3 coding segment. Just upstream, a 13-nucleotide segment matched to a published D segment sequence (Dsp 2.2) (Kurosawa, Y. *et al.*, *J. Exp. Med.*, 155: 201 (1982), and Toneyawa, S., *Nature*, 302: 575 (1983)). A nonapeptide predicted from this area showed characteristic homology to the published mouse heavy chain V subgroups at amino acid residues 86 to 94, comprising the FR3 of heavy chain molecules. Plasmid pJ3-11 represented a rearranged VDJ sequence, and apparently contained the anti-hepatitis V_H sequence produced by the cell line.

[0159] In order to isolate a V_H region cDNA clone that had a BstEII site in the J region, an *AluI* to *Sau96I*, 265 nucleotide long, probe from pJ3-11 was next used to screen the cDNA library generated from the 13-mer BstEII primer. Six positive clones were isolated. The largest, pBs13-1, was further analyzed. The insert was 280 nucleotides long and its restriction map agreed with that of pJ3-11 except for the introduced BstEII site. Figure 9 illustrates how these two inserts were recombined to generate pMVHCa-13, a V_H clone with the module-joining BstEII site. Three additional V_H cDNA clones were isolated from a cDNA library generated from the 21-mer oligonucleotide UIG-MJH3BSTEII primer containing a BstEII site. These clones may provide alternate V_H cDNA sequences to join to human C_H sequences.

(2) Preparation of cDNA Clones, and Vehicles Containing Same, for Light Chain Mouse Anti-HBsAg Variable Region

[0160] Since the J_κ2 sequence is present in mRNA prepared from the anti-hepatitis hybridoma cell line, the oligonucleotide UIG-JK2BGLII (Figure 7B), was designed to introduce a BglII site into the J_κ2 region. Digestion with BglII would then allow direct insertion of a V_κ cDNA coding region into the BglII site of the previously noted human C_κ vector, pJG2001. This insertion would result in the precise joining of a mouse variable region segment (including the J region) to a human *kappa* constant region segment, each in the proper coding frame and with no alteration in amino acid sequence for either mouse variable or human constant region.

[0161] The JK2BGLII oligonucleotide was used to prime anti-HBsAg mRNA to form a cDNA library as for heavy chain, *supra*, in pUC9. The cDNA was size-selected by polyacrylamide gel electrophoresis prior to cloning, and 50% of the cDNA clones were shown to have insert sizes between 300 and 750 nucleotides in length. Replica filters of this library were screened with two oligonucleotides, the original primer and a second probe complementary to J_κ2 sequence 5' to the original primer.

[0162] It was discovered that the anti-hepatitis B monoclonal cell line CRL 8017 secretes immunoglobulins with at least two different light chains. One of them is derived from the myeloma NS-1, which was used as a fusion partner in generating the anti-hepatitis B cell line. Since NS-1 is derived from the myeloma MOPC21, the possibility was investigated that MOPC21 V_κ mRNA may be present in the V_κ cDNA library from the anti-hepatitis monoclonal cell line. Indeed, one cDNA clone (p6D4B) analyzed has an identical restriction enzyme map to that of MOPC21 V_κ cDNA, except for the inserted BglII site.

[0163] Two conclusions can be drawn from these results. The first is that it is possible to effectively use an oligonucleotide to introduce a restriction enzyme site while cloning a V_κ region from a hybridoma cell line. The second is that one must carefully monitor hybridoma cell lines for the presence of multiple V region sequences, only one of which is the desired sequence.

[0164] In order to further characterize the *kappa* light chain J regions present in the cell line mRNA, poly(A)⁺ RNA was bound to nitrocellulose by the formaldehyde "Dot blot" procedure of White and Bancroft, *J. Biol. Chem.*, 257: 8569 (1982). The RNA was hybridized to 32P-labeled oligonucleotide probes specific for each functional *kappa* J region. These probes are shown in Figure 7B as the UIG probes 5JK1, MJK, 5JK4, and 5JK5. The results showed that the mRNA hybridized strongly to both MJK and 5JK4 oligonucleotide probes, indicating that both J_κ2 and J_κ4 sequences were present. Since J_κ2 mRNA had been previously identified as the one derived from the parental hybridoma partner NS-1, it was concluded that the J_κ4 mRNA encoded the anti-hepatitis binding specificity of the CRL 8017 cells.

[0165] Two different cDNA libraries were screened to isolate V region clones encoding J_κ4 sequences. The first was primed by JK2BGLII, *supra*. The second was made by using the oligonucleotide primer, JK4BGLII, which is specific for J_κ4 mRNA and introduces a BglII site into the J region of cloned V regions. The JK4BGLII primer was used to prime first strand cDNA synthesis to construct a cDNA library by the same method used to construct a JK2BGLII primed cDNA library, except that cDNA was not size selected prior to cloning.

[0166] Figure 7B tabulates the mismatches that each primer has with other functional mouse *kappa* J region sequences. Note that J_κ4 has five mismatches in 21 nucleotides when compared with the JK2BGLII primer, and 3 in 23 with the JK4BGLII primer.

[0167] Both libraries were screened for V region clones containing J_κ4 sequences by hybridizing to an oligonucleotide probe specific for J_κ4 sequences (5JK4). The results of this screen are shown in Table 1.

Table 1*

Library	Probe Specificity	
	J _K 2	J _K 4
JK2BGLII	2% (30/1500)	0.15% (2/1500)
JK4BGLII	N/D	3.5% (31/875)

* Percentage of clones containing J_K2 or J_K4 sequence plus a V region. The probes used were the oligonucleotide 5JK4 (J_K4 specificity, Figure 7) and p6D4B, which contains the NS-1 (MOPC21) V region sequence. N/D, not done.

[0168] Several J_K4 V region cDNA clones isolated from both libraries were characterized. These clones have identical restriction enzyme maps, including the engineered BglII site resulting from the oligonucleotide primed cDNA cloning procedure. The restriction map and sequence of one clone, pV17, show that pV17 contains V region gene sequences.

[0169] These results show that the JK2BGLII primer could correctly, although inefficiently, prime J_K4 mRNA sequences. Since the JK2BGLII primer had less mismatches with any other J_K region mRNA than with J_K4 mRNA (Figure 78), it is expected that the other JK mRNAs can be primed at the correct location with better efficiency using the JK2BGLII primer. Thus, efficient cDNA cloning of any functional mouse kappa V region may be obtained by using a mixture of the JK2BGLII and JK4BGLII primers.

[0170] The placement of a BglII site into the J region during cDNA cloning of the V regions allows joining of the cloned mouse V region gene module to the human kappa constant region gene module (Figure 9B).

[0171] After the aforementioned experiments were carried out it was found that the cDNA clone pV17 lacked a complete 5' coding region. Nucleotide sequencing showed that the A of the initiator codon ATG was not cloned in pV17. This was not a random cDNA cloning artifact because two other cDNA clones had the same defect. Two approaches were devised to obtain a light chain gene with a complete 5' coding region.

[0172] First, a new cDNA library was constructed by first priming with an oligonucleotide (5'-ATATTGCT-GATGCT CT-3') complementary to pV17 sequences 155 bases from the 5' end. From this library, clones hybridizing to a pV17 DNA fragment probe were selected, and some of these new cDNA clones have the initiator ATG plus about 20 nucleotides of 5' untranslated region. One of these clones, p2-12, supplies a 5' untranslated region of 23 nucleotides and a complete ATG initiator codon. When p2-12 was combined with pV17 derived sequences, a variable region with a complete 5' end was formed (p1NG2013E).

[0173] Second, site-directed mutagenesis on the existing light chain clone was used to simultaneously remove the poly-G tract and place a ribosome recognition sequence adjacent to the initiator ATG. The PstI fragment from pV17 was subcloned into M13mp18. An oligonucleotide (V17-IVM; 5'-GTGTCGACTCAGCATGAGGTTC AGGTTTC-3') was then used as a primer to mutate the pV17 sequence to include a SalI site and an initiator ATG into the pV17 sequence. The resultant plasmid pV17-IVM provided an alternate mouse variable region for joining to human constant region modules.

[0174] The complete nucleotide sequence of the variable region from pV17 was then determined. The sequence shows that pV17 contains a V_K-J_K junction region, containing several conserved amino acids, and the hybrid J_K2/J_K4 region formed by priming the J_K4 RNA with the UIG-JK2BGLII oligonucleotide. However, the V_K region in pV17 is non-functional, because the V_K and J_K regions are not in the same coding frame. Translation of the pV17 V region would thus result in an abnormal immunoglobulin light chain where the J region is translated in an incorrect frame. This defect may be caused by aberrant V-J joining, resulting in a non-functional kappa mRNA, as has been observed by Kelley, D.E. *et al.*, *Mol. Cell. Biol.*, 5: 1660-1675 (1985).

[0175] Since the pV17 V region encodes an abnormal immunoglobulin, it is highly unlikely that this light chain is part of a functional anti-hepatitis antibody molecule. These results show the importance of monitoring hybridoma cells for the presence of multiple RNA species encoding V regions, only one of which is the desired sequence.

[0176] Further screening of CRL 8017 cDNA libraries was done to search for V_K cDNA clones which are not from either of the two V_K cDNA classes found so far (MOPC21-p6D4B, pV17). First an oligo-dT primed cDNA library made from CRL8017 RNA was screened with a DNA fragment probe specific for the kappa constant region, and separately with probes specific for MOPC21 and pV17 V_K regions. A cDNA clone (pIE9L-81) that contains the kappa constant region, but has a different V_K region than that of MOPC21 or pV17 was discovered. This method of screening oligo-dT primed cDNA libraries is a useful alternative to oligonucleotide screening of cDNA libraries, because nick-translated probes of high specific activity are used. Also, this method allows the simultaneous isolation of several classes of V region clones, such as all V_K clones, by appropriate probe choice. Second, the UIG-JK2BGLII-primed cDNA library made from CRL 8017 RNA was screened with the UIG-5JK2 oligonucleotide probe (see Figure 7). A new class of V_K cDNA clones was found whose members are homologous to pIE9L-81 and hybridize to the UIG-5JK2 probe, but not

to a MOPC21 V_K probe. The restriction endonuclease site maps and nucleotide sequences of these clones also differ from MOPC21-homologous V_K cDNA clones from CRL8017 cells. These clones, however, have an aberrant V-J joint which results in a nonfunctional mRNA, and appear to be identical to one described by Cabilly and Riggs (*Gene*, **40**: 157 (1985)).

[0177] It was therefore concluded that the anti-hepatitis B cell line CRL8017 has at least three classes of V_K mRNA corresponding to the above described cDNA clones p6D4B (MOPC21), pIE9L, and pV17. The pIE9L and pV17 clones are derived from mRNA from aberrantly rearranged Kappa genes, while the p6D4B clone is derived from the parent hybridoma fusion partner NS-1. None of these clones appear to encode the desired anti-hepatitis light chain.

(3) Preparation and Expression of Heavy Chain Containing Human Constant/Mouse Variable Regions

[0178] The V region sequences in pMVHCa-13 were joined to the human IgG1 constant (C) region clone pGMH-6. Due to the presence of a second *Bst*II site within the IgG1 CH1 region of pGMH-6, a multi-step ligation was required. First, the 220 nucleotide *Bst*II fragment from the J-CH1 region of pGMH-6 was ligated to the 1100 nucleotide IgG region *Bst*II to *Bam*HI fragment of pGMH-6. In a separate ligation, the 420 nucleotide *Bst*II to *Bam*HI fragment of pMVHCa-13, which comprises the mouse V region, was joined to a calf intestine phosphatase treated *Bam*HI plasmid vector. The two ligations were then combined, ligase was added, and the products were transformed into HB101, resulting in the chimeric mouse V-human C clone pMVHCc-24 (Figure 9A).

[0179] The V region of the hybrid heavy chain gene in pMVHCc-24 was further analyzed by partial sequence analysis. This analysis showed that the cloned V region contained a D sequence which matches a known D sequence, DSP2.2 (Kurosawa and Tonegawa, *supra*). The sequence also predicted a 19 amino acid leader peptide similar to known mouse V heavy chain leader peptide sequences, and a 5' untranslated region of at least 3 nucleotides.

[0180] The *Bam*HI fragment containing the mouse-human hybrid heavy chain gene of pMVHCc-24 was cloned into *Bam*HI digested pING2003E vector, resulting in the expression plasmid pING2006E (Figure 11). The pING2006E plasmid should have an increased probability of efficient expression of the mouse-human chimeric immunoglobulin gene in B lymphoid cells because of the presence of the mouse heavy chain enhancer region.

[0181] A modification of the chimeric heavy chain gene present in pMVHCc-24 was done to provide an alternate heavy chain gene which lacks the oligo-dC region preceding the initiator ATG. The pING2012E and pING2006E vectors are identical except for the nucleotides immediately preceding the ATG, as shown in Figure 12.

[0182] Bacteria harboring the pING2006E and pSV2-neo plasmids were converted into protoplasts by the method of Sandri-Goldin, R. M. et al., *Mol. Cell. Biol.*, **1**: 743 (1981). The protoplasts were then separately fused to SP2/0-Ag14 hybridoma cells (ATCC CRL 1581) by treatment with polyethyleneglycol (Ochi, A. et al., *Nature*, **302**: 340, 1983). The fused cells were allowed to recover for 72 hours in complete medium before plating at 10,000 or 50,000 cells per well in a 96-well tissue culture plate. The cells were selected with G418 at 0.8 mg/ml for two weeks, when growth in some wells was clearly evident. Under these selection conditions, SP2/0 cells were completely killed within 4-7 days by G418. Only cells which have integrated and expressed the neo gene present in the vectors will grow under G418 selection. The number of wells positive for growth by these integrative transfectants are shown in Table 2.

Table 2*

Strain/Plasmid	10,000 cells/well	50,000 cells/well
MC1061/pING2006E	3 (13%)	12 (50%)
MC1061/pSV2-neo	7 (29%)	4 (17%)
MC1061/none	0	0
* Percentage of wells showing positive growth out of 24 wells.		

[0183] Cells transfected with pING2006E and pSV2-neo were tested for immunoglobulin gene expression at the RNA and protein level. Total cell RNA was prepared from transfected cells, bound to nitrocellulose and hybridized to nick-translated probes specific for the mouse-human hybrid heavy chain gene. Two clones were found which have a strong signal, representing expression of the gene at the RNA level. The amount of total cellular RNA hybridizing to the mouse-human probe appeared to be approximately 1/10 the level of heavy chain RNA in the original hybridoma cells. This probably represented about 1% of the total mRNA of the transfected cell.

[0184] The transfected mouse cells were also tested for production of cytoplasmic human heavy chain protein by an ELISA assay. It was found that 3 out of 7 pING2006E transfected cell lines produced detectable levels of human heavy chain protein. The mouse cell transformant producing the most mouse-human heavy chain protein gave a signal in the ELISA assay comparable to that of a 1/100 dilution of a human B cell line producing intact human immunoglobulin

IgG1. This modest level of detected mouse-human heavy chain protein may be due to several factors, including instability of heavy chains in the absence of light chains in hybridoma cells, or incorrect processing of the chimeric gene transcript.

(4) Gene Amplification of the Integrated Chimeric Gene

[0185] Southern blot analysis showed that multiple copies of the pING2006E DNA sequences were integrated in tandem in the mouse genome. Restriction enzymes ApaI and BglII both cleave pING2006E singly. In the transformant, 2AE9, a band, from an ApaI or BglII digestion, of the expected size (8.2kb) was found to hybridize to the human C gamma 1 sequences (data not shown). An BamHI band of the correct size (1.6kb) was found to hybridize to the human as well as the IE9 V_H sequences. Gene-copy titration experiment (Fig. 14) indicated that there are about 5 copies of pING2006E in a 2AE9 genome. That fact that only a single band was detected in the ApaI or BglII lane indicates that these individual copies are in a tandemly arranged array. A set of double digestions showed that pING2006E sequences suffered no rearrangement in their introduction into the mouse DNA (data not shown).

[0186] We next transfected the 2AE9 cells with a plasmid that contains a different selectable marker, the gpt gene, and selected clones growing out in DMEM-HAT. One clone, 2BH10, has about 38 ng soluble human gamma 1 protein per 10⁶ cells. Southern analysis showed that 2BH10 has about 30 copies of pING2006E (Fig. 14). They were amplified from the 5 copies in 2AE9 without rearrangement of the DNA sequences. (Compare the 2AE9 panel to the 2BH10). SI data (data not shown) revealed that this increase in template led to a higher amount of IgG gene transcripts. We believe that these sequences were co-amplified with contiguous cellular sequences as a result of the second selection.

EXAMPLE III: A Human-Mouse Chimeric Antibody with Cancer Antigen Specificity

(1) Antibody L6

[0187] L6 monoclonal antibody (MAb) was obtained from a mouse which had been immunized with cells from a human lung carcinoma, after which spleen cells were hybridized with NS-1 mouse myeloma cells. The antibody binds to a previously not identified carbohydrate antigen which is expressed in large amounts at the surface of cells from most human carcinomas, including lung carcinomas (adeno, squamous), breast carcinomas, colon carcinomas and ovarian carcinomas, while the antigen is only present at trace levels in normal cells from the adult host. MAb L6 is an IgG2a and can mediate antibody dependent cellular cytotoxicity, ADCC, in the presence of human peripheral blood leukocytes as a source of effector cells, so as to lyse L6 positive tumor cells, and it can lyse L6 positive tumor cells in the presence of human serum as a source of complement; the lysis is detected as the release of ⁵¹Cr from labelled cells over a 4 hour incubation period. MAb L6 can localize to L6 positive tumors xenotransplanted onto nude mice, and it can inhibit the outgrowth of such tumors. MAb L6 is described in Cancer Res. **46**:3917-3923, 1986 (on MAb specificity) and in Proc. Natl. Acad. Sci. **83**:7059-7063, 1986 (on MAb function).

(2) Identification of J Sequences in the Immunoglobulin mRNA of L6.

[0188] Frozen cells were thawed on ice for 10 minutes and then at room temperature. The suspension was diluted with 15 ml PBS and the cells were centrifuged down. They were resuspended, after washes in PBS, in 16 ml 3M LiCl, 6M urea and disrupted in a polytron shear. The preparation of mRNA and the selection of the poly(A)⁺ fraction were carried out according to Auffray, C. and Rougeon, F., Eur. J. Biochem. **107**:303, 1980.

[0189] The poly (A)⁺ RNA from L6 was hybridized individually with labeled J_H1, J_H2, J_H3 and J_H4 oligonucleotides under conditions described by Nobrega et al. Anal. Biochem. **131**:141, 1983). The products were then subjected to electrophoresis in a 1.7% agarose-TBE gel. The gel was fixed in 10% TCA, blotted dry and exposed for autoradiography. The result showed that the L6 V_H contains J_H2 sequences.

[0190] For the analysis of the V_K mRNA, the dot-blot method of White and Bancroft J. Biol. Chem. **257**:8569, (1982) was used. Poly (A)⁺ RNA was immobilized on nitrocellulose filters and was hybridized to labeled probe-oligonucleotides at 40° in 4xSSC. These experiments show that L6 contains J_K5 sequences. A faint hybridization to J_K2 was observed.

(3) V Region cDNA Clones.

[0191] A library primed by oligo (dT) on L6 poly (A)⁺ RNA was screened for kappa clones with a mouse C_K region probe. From the L6 library, several clones were isolated. A second screen with a 5' J_K5 specific probe identified the L6 (J_K5) light-chain clones. Heavy chain clones of L6 were isolated by screening with the J_H2 oligonucleotide.

[0192] The heavy and light chain genes or gene fragments from the cDNA clones, pH3-6a and pL3-12a were inserted into M13 bacteriophage vectors for nucleotide sequence analysis. The complete nucleotide sequences of the variable

region of these clones were determined (FIGURES 15 and 16) by the dideoxy chain termination method. These sequences predict V region amino acid compositions that agree well with the observed compositions, and predict peptide sequences which have been verified by direct amino acid sequencing of portions of the V regions.

[0193] The nucleotide sequences of the cDNA clones show that they are immunoglobulin V region clones as they contain amino acid residues diagnostic of V domains (Kabat *et al.*, *Sequences of Proteins of Immunological Interest*; U.S. Dept. of HHS, 1983).

[0194] The L6 V_H belongs to subgroup II. The cDNA predicts an N-terminal sequence of 24 amino acid residues identical to that of a known V_H (45-165 CFI; Margolies *et al.* *Mol. Immunol.* 18:1065, 1981). The L6 V_H has the J_H2 sequence. The L6 V_L is from the V_K-Kpnl family (Nishi *et al.* *Proc. Nat. Acad. Sci. USA* 82:6399, 1985), and uses J_K5. The cloned L6 V_L predicts an amino acid sequence which was confirmed by amino acid sequencing of peptides from the L6 light chain corresponding to residues 18-40 and 80-96.

(4) In Vitro Mutagenesis to Engineer Restriction Enzyme Sites in the J Region for Joining to a Human C-Module, and to Remove Oligo (dC) Sequences 5' to the V Modules.

[0195] Both clones generated from priming with oligo (dT) L6 V_K and L6 V_H need to be modified. For the L6 V_K, the J-region mutagenesis primer J_KHindIII, as shown in FIGURE 17B, was utilized. A human C_K module derived from a cDNA clone was mutagenized to contain the HindIII sequence (see FIGURE 17A). The mutagenesis reaction was performed on M13 subclones of these genes. The frequency of mutant clones ranged from 0.5 to 1% of the plaques obtained.

[0196] It had been previously observed that the oligo (dC) sequence upstream of the AUG codon in a V_H chimeric gene interferes with proper splicing in one particular gene construct. It was estimated that perhaps as much as 70% of the RNA transcripts had undergone the mis-splicing, wherein a cryptic 3' splice acceptor in the leader sequence was used. Therefore the oligo (dC) sequence upstream of the initiator AUG was removed in all of the clones.

[0197] In one approach, an oligonucleotide was used which contains a Sall restriction site to mutagenize the L6 V_K clone. The primer used for this oligonucleotide-directed mutagenesis is a 22-mer which introduces a Sall site between the oligo (dC) and the initiator met codon (FIGURE 19).

[0198] In a different approach, the nuclease BAL-31 was used to chew away the oligo (dC) in the L6 V_H clone pH3-6a. The size of the deletion in two of the mutants obtained was determined by nucleotide sequencing and is shown in FIGURE 17. In both of these mutants (delta 4 and delta 21), all of the oligo (dC) 5' to the coding region were deleted.

[0199] These clones were then modified by oligonucleotide-directed mutagenesis with the MJH2-ApaI primer (FIGURE 17). This 31-base primer introduces an ApaI site in the mouse C_H gene at a position analogous to an existing ApaI site in human C γ 1 cDNA gene module. The primer introduces the appropriate codons for the human C γ 1 gene. The chimeric heavy chain gene made by joining the mutagenized mouse V_H gene module to a human C_H module thus encodes a chimeric protein which contains no human amino acids for the entire V_H region.

[0200] The human C γ 1 gene module is a cDNA derived from GM2146 cells (Human Genetic Mutant Cell Repository, Newark, New Jersey). This C γ 1 gene module was previously combined with a mouse V_H gene module to form the chimeric expression plasmid pING2012E.

(5) L6 Chimeric Expression Plasmids.

[0201] L6 chimeric heavy chain expression plasmids were derived from the replacement of the V_H module pING2012E with the V_H modules of mutants delta 21 and delta 4 to give the expression plasmids pING2111 and pING2112 (FIGURE 17). These plasmids direct the synthesis of chimeric L6 heavy chain when transfected into mammalian cells.

[0202] For the L6 light chain chimeric gene, the Sall to HindIII fragment of the mouse V_K module was joined to the human C_K module by the procedure outlined in FIGURE 18, forming pING2119. Replacement of the neo sequence with the *E. coli* gpt gene derived from pSV2-gpt resulted in pING2120, which expressed L6 chimeric light chain and confers mycophenolic acid resistance when transfected into mammalian cells.

[0203] The inclusion of both heavy and light chain chimeric genes in the same plasmid allows for the introduction into transfected cells of a 1:1 gene ratio of heavy and light chain genes leading to a balanced gene dosage. This may improve expression and decrease manipulations of transfected cells for optimal chimeric antibody expression. For this purpose, the DNA fragments derived from the chimeric heavy and light chain genes of pING2111 and pING2119 were combined into the expression plasmid pING2114 (FIGURE 19). This expression plasmid contains a selectable neo^R marker and separate transcription units for each chimeric gene, each including a mouse heavy chain enhancer.

[0204] The modifications and V-C joint regions of the L6 chimeric genes are summarized in FIGURE 20.

(6) Stable Transfection of Mouse Lymphoid Cells for the Production of Chimeric Antibody.

[0205] Electroporation was used (Potter *et al.* supra, Toneguzzo *et al.* Mol. Cell Biol. 6:703 1986) for the introduction of L6 chimeric expression plasmid DNA into mouse Sp2/0 cells. The electroporation technique gave a transfection frequency of $1 \cdot 10 \times 10^{-6}$ for the Sp2/0 cells.

[0206] The two gene expression plasmid pING2114 was linearized by digestion with *Aat*II restriction endonuclease and transfected into Sp2/0 cells, giving approximately fifty G418 resistant clones which were screened for human heavy and light chain synthesis. The levels of chimeric antibody chain synthesis from the two producers, D7 and 3E3, are shown in Table 3. Chimeric L6 antibody was prepared by culturing the D7 transfectant cells for 24 hours at 2×10^6 cells/ml in 5 l DMEM supplemented with HEPES buffer and penicillin and streptomycin. The supernatant was concentrated over an Amicon YM30 membrane in 10mM sodium phosphate buffer, pH8.0. The preparation was loaded over a DEAE-Cellulose column, which separated the immunoglobulin into unbound and bound fractions. Samples from the DEAE-unbound, DEAE-bound and the pre-DEAE preparations (from 1.6 ul of medium) was separately purified by affinity chromatography on a Protein-A Sepharose column, eluting with 0.1 M sodium citrate, pH 3.5. The eluted antibody was neutralized and concentrated by Amicon centricon filtration, in phosphate-buffered saline. The yields for the three preparations were 12ug (DEAE unbound), 6ug (DEAE bound), and 9ug (pre-DEAE column). Western analysis of the antibody chains indicated that they were combined in an H_2L_2 tetramer like native immunoglobulins.

(7) A second purification for Chimeric L6 Antibody Secreted in Tissue Culture.

[0207]

- a. Sp2/0.pING2114.D7 cells were grown in culture medium [DMEM (Gibco #320-1965), supplemented with 10% Fetal Bovine Serum (Hyclone #A-1111-D), 10mM HEPES, 1x Glutamine-Pen-Strep (Irvine Scientific #9316) to 1×10^6 cell/ml.
- b. The cells were then centrifuged at 400xg and resuspended in serum-free culture medium at 2×10^6 cell/ml for 18-24 hr.
- c. The medium was centrifuged at 4000 RPM in a JS-4.2 rotor (3000xg) for 15 min.
- d. 1.6 liter of supernatant was then filtered through a 0.45 micron filter and then concentrated over a YM30 (Amicon Corp.) filter to 25ml.
- e. The conductance of the concentrated supernatant was adjusted to 5.7-5.6 mS/cm and the pH was adjusted to 8.0.
- f. The supernatant was centrifuged at 2000xg, 5 min., and then loaded onto a 40 ml DEAE column, which was pre-equilibrated with 10mM sodium phosphate, pH8.0.
- g. The flow through fraction was collected and loaded onto a 1ml protein A-Sepharose (Sigma) column pre-equilibrated with 10mM sodium phosphate, pH8.0.
- h. The column was washed first with 6ml 10mM sodium phosphate buffer pH=8.0, followed by 8ml 0.1M sodium citrate pH=3.5, then by 6ml 0.1M citric acid (pH=2.2). Fractions of 0.5ml were collected in tubes containing 50ul 2M Tris base (Sigma).
- i. The bulk of the IgG was in the pH=3.5 elution and was pooled and concentrated over Centricon 30 (Amicon Corp.) to approximately .06ml.
- j. The buffer was changed to PBS (10mM sodium phosphate pH=7.4, 0.15M NaCl) in Centricon 30 by repeated diluting with PBS and re-concentrating.
- k. The IgG solution was then adjusted to 0.10ml and bovine serum albumin (Fraction V, U.S. Biochemicals) was added to 1.0% as a stabilizing reagent.

(8) Production and Purification of Chimeric L6 Antibody Secreted in Ascites Fluid.

[0208]

- a. The ascites was first centrifuged a 2,000 xg for 10 min.
- b. The conductance of the supernatant was adjusted to 5.7-5.6 mS/cm and its pH adjusted to 8.0.
- c. Supernatant was then loaded onto a 40 ml DEAE-cellulose column pre-equilibrated with 10 mM Na_2HPO_4 pH8.0.
- d. The flow through from the DEAE column was collected and its pH was adjusted to 7.4, and then loaded onto a 1.0 ml goat anti-human IgG (H + L) - sepharose column.
- e. The column was washed first with 6 ml of 10 mM sodium phosphate, 0.5 M sodium chloride, followed by 8 ml of 0.5 M NH_4OH , and 3 M sodium thiocyanate.
- f. The sodium thiocyanate eluate was pooled and dialyzed against 2L PBS overnight.

[0209] The antibody can be further concentrated by steps j. and k. of the previous procedure.

TABLE 3

Levels of Secreted Chimeric L6 Chains from Sp2/0 Transfectants*					
Culture Condition	FBS	Sp2/0.D7		Sp2/0.3E3	
		Kappa ^b	Gamma ^c	Kappa ^b	Gamma ^c
1. 20 ml, 2d, seed @ 2x10 ⁶ /ml	+	17	77	100	700
2. 200 ml, 2d, seed @ 2.5x10 ⁶ /ml	+	0.9	6	80	215
3. 200 ml, 1d, seed @ 2x10 ⁶ /ml	-	1.9	3.8	97	221
4. Balb/c ascites	-	5,160	19,170	ND	ND
ND - Not determined.					
FBS: Fetal Bovine Serum					

* - Sp2/0 cells transfected by electroporation with pING2114(pL6HL).

^b - ug/l measured by ELISA specific for human Kappa - human Bence-Jones protein standard.

^c - ug/l measured by ELISA specific for human gamma - human IgG standard.

(9) Studies Performed on the Chimeric L6 Antibody.

[0210] First, the samples were tested with a binding assay, in which cells of both an L6 antigen-positive and an L6 antigen-negative cell line were incubated with standard mouse monoclonal antibody L6, chimeric L6 antibody derived from the cell culture supernatants, and chimeric L6 antibody derived from ascites (as previously described) followed by a second reagent, fluorescein-isothiocyanate (FITC)-conjugated goat antibodies to human (or mouse, for the standard) immunoglobulin.

[0211] Since the binding assay showed strong reactivity of the chimeric L6 on the L6 antigen positive cell line and total lack of reactivity on the negative cell line, the next step was to test for the ability of the chimeric L6 to inhibit the binding of mouse L6 to antigen positive cells; such inhibition assays are used routinely to establish the identity of two antibodies' recognition of antigen. These data are discussed below ("Inhibition of binding"). As part of these studies, a rough estimate of antibody avidity was made.

[0212] Finally, two aspects of antibody function were studied, the ability to mediate ADCC in the presence of human peripheral blood leukocytes, and the ability to kill L6 positive tumor cells in the presence of human serum as a source of complement (see "Functional Assays" below).

[0213] Binding Assays. Cells from a human colon carcinoma line, 3347, which had been previously shown to express approximately 5 x 10⁶ molecules of the L6 antigen at the cell surface, were used as targets. Cells from the T cell line HSB2 was used as a negative control, since they, according to previous testing, do not express detectable amounts of the L6 antigen. The target cells were first incubated for 30 min at 4°C with either the chimeric L6 or with mouse L6 standard, which had been purified from mouse ascites. This was followed by incubation with a second, FITC-labelled, reagent, which for the chimeric antibody was goat-anti-human immunoglobulin, obtained from TAGO (Burlingame, CA), and used at a dilution of 1:50. For the mouse standard, it was goat-anti-mouse immunoglobulin, also obtained from TAGO and used at a dilution of 1:50. Antibody binding to the cell surface was determined using a Coulter Model EPIC-C cell sorter.

[0214] As shown in Table 4 and Table 4A, both the chimeric and the mouse standard L6 bound significantly, and to approximately the same extent, to the L6 positive 3341 line. They did not bind above background to the L6 negative HSB2 line.

[0215] In view of the fact that the three different chimeric L6 samples presented in Table 4 behaved similarly in the binding assays, they were pooled for the inhibition studies presented below. The same inhibition studies were performed for chimeric L6 derived from ascites fluid presented in Table 4A.

[0216] Inhibition of Binding. As the next step was studied the extent to which graded doses of the chimeric L6 antibody, or the standard mouse L6, could inhibit the binding of an FITC-labelled mouse L6 to the surface of antigen positive 3347 colon carcinoma cells.

[0217] Both the chimeric and mouse standard L6 inhibited the binding of the directly labelled L6 antibody, with the binding curves being parallel. The chimeric antibody was slightly less effective than the standard, as indicated by the results which showed that 3.4 ug/ml of the pooled chimeric L6 MAb, as compared to 2.0 ug/ml of the standard mouse L6 MAb was needed for 50% inhibition of the binding, and that 5.5 ug/ml of the chimeric L6 (derived from ascites) as compared to 2.7 ug/ml of the standard mouse L6 MAb was needed for 50% inhibition of binding.

[0218] As part of these studies, a rough estimate was made of antibody avidity. The avidity of the standard mouse

L6 had been previously determined to be approximately 4×10^6 . The data indicated that there were no significant differences in avidity between the chimeric and the mouse L6.

[0219] Functional Assays. A comparison was made between the ability of the chimeric L6 and standard mouse L6 to lyse L6 antigen positive cells in the presence of human peripheral blood leukocytes as a source of effector cells (mediating Antibody Dependent Cellular Cytotoxicity, ADCC) or human serum as a source of complement (mediating Complement-Dependent Cytolysis, CDC).

[0220] As shown in Table 5 and Tables 5A-5D, the chimeric L6 was superior to the simultaneously tested sample of mouse L6 in causing ADCC, as measured by a 4 hr ^{51}Cr release test.

[0221] Tables 6 and 6A-6B present the data from studies on complement-mediated target cell lysis. In this case, a high cytolytic activity was observed with both the mouse and the chimeric L6 antibodies.

Conclusions

[0222] The results presented above demonstrate a number of important, unexpected qualities of the chimeric L6 monoclonal antibody of the invention. Firstly, the chimeric L6 antibody binds to L6 antigen positive tumor cells to approximately the same extent as the mouse L6 standard and with approximately the same avidity. This is significant for the following reasons: the L6 antibody defines (a) a surface carbohydrate antigen, and (b) a protein antigen of about 20,000 daltons, each of which is characteristic of non-small cell lung carcinoma (NSCLC) and certain other human carcinomas. Significantly, the L6 antibody does not bind detectably to normal cells such as fibroblasts, endothelial cells, or epithelial cells in the major organs. Thus the chimeric L6 monoclonal antibody defines an antigen that is specific for carcinoma cells and not normal cells.

[0223] In addition to the ability of the chimeric L6 monoclonal antibodies of the present invention to bind specifically to malignant cells and localize tumors, the chimeric L6 exerts profound biological effects upon binding to its target, which make the chimeric antibody a prime candidate for tumor immunotherapy. The results presented herein demonstrate that chimeric L6 is capable of binding to tumor cells and upon binding kills the tumor cells, either by ADCC or CDC. Such tumor killing activity was demonstrated using concentrations of chimeric L6 antibody as low as 0.01 $\mu\text{g/ml}$ (10ng/ml).

[0224] Although the prospect of attempting tumor therapy using monoclonal antibodies is attractive, with some partial tumor regressions being reported, to date such monoclonal antibody therapy has been met with limited success (Houghton, February 1985, Proc. Natl. Acad. Sci. 82:1242-1246). The therapeutic efficacy of mouse monoclonal antibodies (which are the ones that have been tried so far) appears to be too low for most practical purposes. The discovery of the profound biological activity of chimeric L6 coupled with its specificity for a carcinoma antigen makes the chimeric L6 antibody a choice therapeutic agent for the treatment of tumors *in vivo*. Moreover, because of the "human" properties which will make the chimeric L6 monoclonal antibodies more resistant to clearance *in vivo*, the chimeric L6 monoclonal antibodies will be advantageously used not only for therapy with unmodified chimeric antibodies, but also for development of various immunoconjugates with drugs, toxins, immunomodulators, isotopes, etc., as well as for diagnostic purposes such as *in vivo* imaging of tumors using appropriately labelled chimeric L6 antibodies. Such immunoconjugation techniques are known to those skilled in the art and can be used to modify the chimeric L6 antibody molecules of the present invention.

[0225] Two illustrative cell lines secreting chimeric L6 antibody were deposited prior to the filing date of this application at the ATCC, Rockville Maryland. These are transfected hybridoma C255 (corresponds to 3E3 cells, supra), ATCC HB 9240 and transfected hybridoma C256 (C7 cells, supra), ATCC HB 9241.

[0226] The present invention is not to be limited in scope by the cell lines deposited since the deposited embodiment is intended as a single illustration of one aspect of the invention and all cell lines which are functionally equivalent are within the scope of the invention. Indeed, various modifications of the invention in addition to those shown in the art from the foregoing description and accompanying drawings are intended to fall within the scope of the appended claims.

TABLE 4

Binding Assays of Chimeric L6 Antibody and Mouse L6 Monoclonal Antibody on an L6 Antigen Positive and L6 Antigen Negative Cell Line.

Antibody	Batch	Binding Ratio For* H3347 Cells (L6 +)	
		GAM	GAH
Standard L6		56.6	4.2
Chimeric L6	a	1.3	110.3
	b	1.3	110.3
	c	1.3	110.3
		Binding Ratio For* HSB-2 Cells (L6 -)	
		GAM	GAH
Standard L6		1.1	1.1
Chimeric L6	a	1.0	1.0
	b	1.0	1.1
	c	1.0	1.1

* All assays were conducted using an antibody concentration of 10 ug/ml. The binding ratio is the number of times brighter a test sample is than a control sample treated with GAM (FITC conjugated goat-anti-mouse) or GAH (FITC conjugated goat anti-human) alone. A ratio of 1 means that the test sample is just as bright as the control; a ratio of 2 means the test sample is twice as bright as the control, etc.

TABLE 4A

Binding Assays Of Chimeric L6 Antibody and Mouse Monoclonal Antibody on an L6 Antigen Positive and L6 Antigen Negative Cell Line.

Antibody	Antibody Concentration (ug/ml)	Binding Ratio For* H3347 Cells (L6 +)	
		GAM	GAH
Standard L6	30	38	4
	10	49	4
	3	40	3
Chimeric L6 (Ascites)	30	2	106
	10	2	106
	3	1	42
Chimeric L6 (Cell culture)	30	1	105
	10	1	86
	3	1	44
		Binding Ratio For** HSB-2 Cells (L6 -)	
		GAM	GAH
Standard L6	10	1	1
Chimeric L6 (Ascites)	10	1	1

* The binding ratio is the number of times brighter a test sample is than a control sample treated with GAM (FITC conjugated goat anti-mouse) alone. A ratio of 1 means that the test sample is just as bright as the control; a ratio of 2 means the test sample is twice as bright as the control, etc.

TABLE 4A (continued)

Binding Assays Of Chimeric L6 Antibody and Mouse Monoclonal Antibody on an L6 Antigen Positive and L6 Antigen Negative Cell Line.			
	Antibody Concentration	Binding Ratio For* H3347 Cells (L6 +)	
		Binding Ratio For** HSB-2 Cells (L6 -)	
		GAM	GAH
Chimeric L6 (Cell Culture)	10	1	1

* The binding ratio is the number of times brighter a test sample is than a control sample treated with GAM (FITC conjugated goat anti-human) alone. A ratio of 1 means that the test sample is just as bright as the control; a ratio of 2 means the test sample is twice as bright as the control, etc.

TABLE 5

ADCC of Chimeric L6 (Mouse) L6 Antibodies On Colon Carcinoma Cell Line 3347.			
Antibody	Antibody Concentration (ug/ml)	PBL per Target Cell	% Cytolysis*
Chimeric L6	10	100	64
	5	100	70
	10	0	2
Standard L6	10	100	24
	5	100	17
	10	0	2
None	0	100	1

* The target cells had been labelled with ^{51}Cr and were exposed for 4 hours to a combination of MAbs and human peripheral blood leukocytes (PBL), and the release of ^{51}Cr was measured subsequently. The release of ^{51}Cr (after corrections of values for spontaneous release from untreated cells) is a measure of the percent cytotoxicity.

TABLE 5A

ADCC of Chimeric L6 and Standard (Mouse) L6 Antibodies On Colon Carcinoma Cell Line 3347.			
Antibody	Antibody Concentration (ug/ml)	PBL per Target Cell	% Cytolysis*
Chimeric L6 (Ascites)	20	100	80
	10	100	74
	5	100	71
	2.5	100	71
	20	0	0
Chimeric L6 (Cell Culture)	10	100	84
	5	100	74
	2.5	100	67
	10	0	3
Standard L6	20	100	32
	10	100	26
	20	0	0

* The target cells had been labelled with ^{51}Cr and were exposed for 4 hours to a combination of MAbs and human peripheral blood leukocytes (PBL), and the release of ^{51}Cr was measured subsequently. The release of ^{51}Cr (after corrections of values for spontaneous release from untreated cells) is a measure of the percent cytotoxicity.

TABLE 5B

ADCC of Chimeric L6 and Standard (Mouse) L6 Antibodies On Colon Carcinoma Cell Line 3347.

Antibody	Antibody Concentration (ug/ml)	PBL per Target Cell	% Cytolysis*
Chimeric L6 (Ascites)	5	100	84
	2.5	100	78
	1.25	100	85
	0.63	100	81
	0.31	100	80
	0.16	100	71
	0.08	100	65
	5	0	0
Standard L6	5	100	32
	5	0	0
None	0	100	19

* The target cells had been labelled with ^{51}Cr and were exposed for 4 hours to a combination of MAbs and human peripheral blood leukocytes (PBL), and the release of ^{51}Cr was measured subsequently. The release of ^{51}Cr (after corrections of values for spontaneous release from untreated cells) is a measure of the percent cytolysis.

TABLE 5C

ADCC of Chimeric L6 and Standard (Mouse) L6 Antibodies On Lung Carcinoma Cell Line H2669.

Antibody	Antibody Concentration (ug/ml)	PBL per Target Cell	% Cytolysis*
Chimeric L6 (Ascites)	10	100	35
	1	100	31
	0.1	100	27
	0.01	100	15
	0.001	100	13
	0.0001	0	15
	0.0001	0	15
Standard L6	10	100	9
	1	100	15
None	0	100	9
Chimeric L6 (Ascites)	10	10	19
	1	10	15
	0.1	10	11
	0.01	10	13
	0.001	10	22
	0.0001	10	11
	0.0001	10	11
	0.0001	10	11
Standard L6	10	10	7
	1	10	6

* The target cells had been labelled with ^{51}Cr and were exposed for 4 hours to a combination of MAbs and human peripheral blood leukocytes (PBL), and the release of ^{51}Cr was measured subsequently. The release of ^{51}Cr (after corrections of values for Spontaneous release from untreated cells) is a measure of the percent cytolysis.

TABLE 5C (continued)

ADCC of Chimeric L6 and Standard (Mouse) L6 Antibodies On Lung Carcinoma Cell Line H2669.			
Antibody	Antibody Concentration (ug/ml)	PBL per Target Cell	% Cytolysis*
None	0	10	8
Chimeric L6 (Ascites)	10	0	4
Standard L6	10	0	9

* The target cells had been labelled with ^{51}Cr and were exposed for 4 hours to a combination of MAbs and Human peripheral blood leukocytes (PBL), and the release of ^{51}Cr was measured subsequently. The release of ^{51}Cr (after corrections of values for Spontaneous release from untreated cells) is a measure of the percent cytolysis.

TABLE 5D

ADCC of Chimeric L6 and Standard (Mouse) L6 Antibodies On Colon Carcinoma Cell Line H3347.			
Antibody	Antibody Concentration (ug/ml)	PBL per Target Cell	% Cytolysis*
Chimeric L6 (Ascites)	10	100	62
	1	100	66
	0.1	100	69
	0.01	100	26
	0.001	100	8
	0.0001	0	3
	10	0	0
Standard L6	10	100	19
	1	100	24
		0	0
None	0	100	8

* The target cells had been labelled with ^{51}Cr and were exposed for 4 hours to a combination of MAbs and Human peripheral blood leukocytes (PBL), and the release of ^{51}Cr (after corrections of values for spontaneous release from untreated cells) is a measure of the percent cytolysis.

TABLE 6

Complement-dependent cytotoxic effect of chimeric and standard (mouse) L6 on colon carcinoma cells from line 3347, as measured by a 4-hr ^{51}Cr -release assay. Human serum from a healthy subject was used as the source of complement.		
Antibody	Human complement	% Cytolysis
L6 Standard 10 ug/ml	Yes	90
L6 chimeric 10 ug/ml	Yes	89
L6 Standard 10 ug/ml	No	0
L6 chimeric 10 ug/ml	No	1

TABLE 6A

Complement Dependent Cytotoxic Effect of Chimeric L6 and Standard(Mouse) L6 Antibodies on Colon Carcinoma Cell Line 3347

Antibody	Antibody Concentration (ug/ml)	PBL per Target Cell	% Cytolysis
Chimeric L6 (Ascites)	20	+	29
	10	+	23
	5	+	18
	2.5	+	8
	20	Inactivated	0
	10	0	0
Chimeric L6 (Cell Culture))	20	+	29
	5	+	26
	2.5	+	18
	20	+	4
	10	0	4
Standard L6	20	+	55
	10	+	37
	20	Inactivated	0
	20	0	1
None	0	+	0

* Complement mediated cytotoxicity was measured by a 4 hour ⁵¹Cr-release assay. Human serum from a healthy subject was used as the source of complement.

TABLE 6B

Complement Dependent Cytotoxic Effect of Chimeric L6 and Standard (Mouse) L6 Antibodies on Colon Carcinoma Cell Line 3347

Antibody	Antibody Concentration (ug/ml)	PBL per Target Cell	% Cytolysis*
Chimeric L6 (Ascites)	10	+	209
	5	+	155
	2.5	+	166
	1.25	+	114
	0.6	+	63
	0.3	+	17
	10	0	0
Standard L6	10	+	96
	5	+	83
	2.5	+	48
	1.25	+	18
	0.6	+	7
	0.3	+	4
	10	0	2
None	0	+	0

* Complement mediated cytotoxicity was measured by a 4 hour ⁵¹Cr-release assay. Human serum from healthy subject was used as the source of complement.

EXAMPLE IV: A Human-Mouse Chimeric Antibody with Specificity for Human B-Cell Antigen

[0227] The 2H7 mouse monoclonal antibody (gamma 2bK) recognizes a human B-cell surface antigen, Bp35 (Clark, E. A., *et al.*, *Proc. Natl. Acad. Sci. USA* **82**:1766 (1985)). The Bp35 molecule plays a role in B-cell activation. mRNA was prepared from the 2H7 cell line. Two cDNA libraries were generated - one using the heavy chain UIG-H primer and the other, oligo(dT). One V_H clone, pH2-11, was isolated upon screening with the same UIG-H oligonucleotide. To isolate the light chain clone, a mouse κ -specific DNA fragment was used to screen the oligo(dT) library. Candidate clones were further screened with a mouse J_K sequences. One V_K clone, pL2-12, was thus isolated. The light chain UIG-K was then used to engineer a restriction enzyme site in the J region.

[0228] The two cDNA clones were also modified at the 5' and to remove the artificial oligo d(C) sequence. In pH2-11 this was carried out by using the restriction enzyme *NcoI* which cuts one nucleotide residue 5' of the ATG initiator codon. In pL2-12 this was achieved by an oligonucleotide *in vitro* mutagenesis using a 22-mer containing a *Sall* site.

[0229] The DNA sequences of these two clones are shown in Figures 21, 22. To construct the chimeric heavy chain plasmid, the V_H module was joined to the human C gamma 1 module (pGMH6) at the J_H *BstEII* site, and the chimeric light chain the V_K module was joined to the human C $_K$ module (pGML60) at the J_K *HindIII* site. The expression vector sequences were derived from pING2012-neo as well as pING2016-gpt. The constructed plasmids are pING2101 (V_H C gamma 1-neo), pING2106 (V_K C $_K$ -neo), pING2101 (V_K C $_K$ -gpt), pING2101 and pING2106 were also used to generate plasmids containing both genes. They are pHL2-11 and pHL2-26. In addition, pING2106 and pING2014 were combined to a two light chain plasmid, pLL2-25, to compensate for the poorer (compared to heavy chain) steady-state accumulation of light chain protein in transfected cells. (See Fig. 23.) Fig. 24 shows the changes made to the variable region sequences during the construction.

[0230] The plasmid, pHL2-11, was linearized by *AatII*; and the DNA was used to transfect Sp2/0 cells by electroporation. Transformants were selected in G418-DMEM. One transformant, 1C9, produces 9.3 ng/ml chimeric κ and 33-72 ng/ml chimeric γ 1 protein as assayed by ELISA. Southern analysis of 1C9 DNA showed that there is one copy of the plasmid integrated in Sp2/0 genome.

EXAMPLE V: Secretion of a Functional Chimeric Antibody from Yeast

(1) Fusion of mature chimeric L6 light chain and heavy chain genes to the yeast invertase signal sequence and shortened phosphoglycerate kinase (PGK promoter).

[0231] Yeast cells are capable of recognizing mammalian secretion signal sequences and of directing secretion of mammalian proteins (Hitzman *et al.*, *supra*). There is, however, evidence which suggests that certain native yeast signal sequences are more effective than mammalian signal sequences at directing secretion of some mammalian proteins from yeast (Smith *et al.*, *Science* **229**:1219 (1985)). One example is the signal sequence for the yeast invertase gene. To improve the efficiency of light and heavy chain secretion, the mature light chain and heavy chain sequences were fused to the yeast invertase signal sequence and placed under transcriptional control of the shortened PGK promoter (U.S. Patent Application 797,477) using the strategies outlined in Figures 25 and 26, respectively. An important element of these constructions is the use of *in vitro* mutagenesis to introduce a restriction site at the signal sequence processing site for both the invertase signal sequence (see U.S. Patent Application 797,477) and the light and heavy chain genes. These restriction sites are positioned such that a blunt-ended ligation of restriction enzyme-digested, T-4 DNA polymerase-treated DNA results in in-phase translational fusions of the 5' end of the mature immunoglobulin chains with the 3' end of the yeast invertase signal sequence. Such genes, when expressed in a yeast cell, may direct the synthesis, processing, and secretion of chimeric light and heavy chains with the same primary peptide sequence as chimeric light and heavy chains secreted from transfected mouse Sp2/0 cells. The DNA sequences of the mutagenesis primers used for light and heavy chain genes as well as the corresponding unmutagenized sequences are shown in Figures 25B and 26B, respectively. Using this approach, the L6 chimeric light and heavy chains were fused to the yeast invertase signal sequence and shortened PGK promoter, resulting in plasmids pING1407-7 and pING1415 (Figures 25C and 26C).

(2) Removal of non-yeast 3' untranslated DNA.

[0232] Recent studies on expression of hepatitis B surface antigen in yeast demonstrated that removal of non-yeast 3' and 5' untranslated sequences can result in increased levels of heterologous gene expression in yeast (Kriesnick *et al.*, *Gene* **46**:135 (1986)). The light chain gene sequence of chimeric L6 antibody in pING1407-7 (Figure 25C) contains approximately 200 bp of 3' untranslated DNA followed by 70 bp of poly A and 20 bp of poly G sequences. An initial treatment of the chimeric L6 light chain DNA with the double-stranded exonuclease *Bal31*, removed the poly A and poly G sequences and all but 90 bp of 3' untranslated DNA, generating the plasmid pING2121b (Figure 27). A restriction

fragment from pING2121b containing only C_{α} was cloned into a derivative of pBR322, generating pING1419 (Figure 27). A second *Bal*31 digestion was next used to remove all but 13 bp of non-yeast 3' untranslated DNA generating the plasmid, pING1431 (Figure 27). The chimeric L6 heavy chain gene in pING1415 (Figure 26) also contains extensive 3' untranslated sequence which includes 80 bp of poly A. All but 11 bp of the 3' untranslated DNA were removed using the strategy shown in Figure 28, generating the plasmid pING1429.

[0233] Site-directed *in vitro* mutagenesis can introduce, at a low frequency, unwanted base pair changes in regions of the DNA outside of the area being mutagenized. To ensure that such mutations were not present in the chimeric L6 light and heavy chain sequences which had been cloned into M13 and subjected to site-directed mutagenesis, we constructed light and heavy chain genes fused to the invertase signal sequence and the shortened PGK promoter which consisted of coding sequences that were either confirmed by DNA sequence analysis or proven to be functional by virtue of their expression in transfected mouse Sp2/O cells to produce functional chimeric L6 antibody. The plasmids, pING1439 (light chain, Figure 27) and pING1436 (heavy chain, Figure 28) were generated by these constructions.

(3) Construction of yeast expression plasmids containing chimeric L6 light and heavy chain genes from pING1439 and pING1436, respectively, fused to the PGK polyadenylation signal.

[0234] In order for yeast to produce an intact functional antibody molecule, a balanced synthesis of both light and heavy chain protein within the host cell is preferred. One approach is to place the light and heavy chain genes on separate expression vectors each containing a different selective marker. A yeast strain defective in the selective markers found on the plasmids can then be either simultaneously or sequentially transformed with these plasmids.

[0235] The chimeric L6 light and heavy chain genes from pING1439 (Figure 27) and pING1436 (Figure 28) were cloned as *Bam*HI-*Xho*I and *Bam*HI-*Xho*I fragments, respectively, in two different medium copy number (about 20 copies/cell) expression vectors (yeast-*E. coli* shuttle). One of these, pING804CVS, contains the complete yeast 2-micron circle, the PGK transcription termination and polyadenylation signals, and the *leu2* gene as the selective marker. The other vector, pING1150, contains the yeast origin of replication, *oriY*, a *cis*-acting sequence (REP3) from the yeast endogenous 2-micron plasmid, the PGK transcription termination and polyadenylation signals, and the *ura3* gene as the selective marker. Both plasmids also contain the β -lactamase gene (*bla*) for ampicillin resistance and the bacterial origin of replication (*oriB*) from pBR322 for selection and amplification in bacteria. Four plasmids resulted from these constructions: pING1441--light chain, *leu2* and pING1443--light chain, *ura3* (Figure 29); pING1440--heavy chain, *leu2* and pING1442--heavy chain, *ura3* (Figure 30).

(4) Secretion of chimeric L6 antibody from transformed yeast cells.

[0236] Two separate transformation experiments were performed in an attempt to obtain both light and heavy chain synthesis in yeast cells. Four μ g each of pING1440 and pING1443, and separately of pING1442 and pING1441 were cotransformed into *Saccharomyces cerevisiae* strains BB331C (MATa, *ura3*, *leu2*) by selecting for growth on SD agar (2% glucose, 0.67% yeast-nitrogen base, 2% agar). Ura⁺ Leu⁺ transformants appeared at 2-3 days of incubation at 30°C. Approximately 100 transformants were obtained for pING1440 plus pING1443; only 15 transformants were obtained for pING1442 plus pING1441. Ten colonies were inoculated from each plate into 5 ml SD broth supplemented with 50 mM sodium succinate, pH 5.5, and grown for 65 hours at 30°C. The cells were removed by centrifugation and the culture supernatants analyzed by ELISA for the levels of light chain and heavy chain and for the degree of association of the secreted light and heavy chains. The latter was assessed using a goat anti-human kappa antiserum to coat the micro-titer wells and a peroxidase-labeled goat anti-human gamma antiserum to detect the level of heavy chain bound to the anti-kappa coat. The results of these assays (Table 7) revealed that all of the culture supernatants from the cells transformed with pING1440 (heavy chain, *leu2*) plus pING1443 (light chain, *ura3*) contained a disproportionately high level of light chain protein relative to the levels of heavy chain protein, and no evidence (at least as determined by ELISA) of assembled light and heavy chains. On the other hand, the supernatants from the cells transformed with pING1442 (heavy chain, *ura3*) + pING1441 (light chain, *leu2*) contained a more balanced production of light and heavy chain proteins, and eight of ten isolates appeared to contain some assembled light and heavy chains as determined by ELISA. Two of these isolates, No. 1 and No. 5, produced a significant proportion of assembled light and heavy chain.

TABLE 7

LEVELS OF SECRETED CHIMERIC L6 LIGHT AND HEAVY CHAIN BY YEAST TRANSFORMANTS ^a				
Plasmids ^b	Isolate No.	Kappa ^c	Gamma ^d	Kappa/Gamma ^e
pING1440 + pING1443	1	284	39	0
	2	324	33	0
	3	473	52	0
	4	387	40	0
	5	316	34	0
	6	188	28	0
	7	381	45	0
	8	455	45	0
	9	380	26	0
	10	579	32	0
pING1441 + pING1442	1	128	79	35
	2	150	30	1
	3	124	29	0
	4	185	55	5
	5	114	52	35
	6	139	23	5
	7	149	34	5
	8	245	57	12
	9	202	26	11
	10	157	19	7

a. *S. cerevisiae* strain BB331C (MATa, leu2, ura3) transformed to Ura⁺ Leu⁺ with plasmide carrying ura3 and leu2 with light or heavy chains.

b. Plasmids:

pING1440 = heavy chain + leu2;

pING1443 = light chain + ura3;

pING1442 = heavy chain + ura3;

pING1441 = light chain + leu2.

c. ng/ml measured by ELISA specific for human kappa with human Bence Jones protein as standard.

d. ng/ml measured by ELISA specific for human gamma with human α IgG standard.

e. ng/ml measured by ELISA using anti-human kappa as coating antibody and anti-human gamma as second antibody with human IgG standard.

[0237] Further analysis was performed to determine if this association was the result of the synthesis of an H₂L₂-size protein. The culture supernatants from isolates Nos. 1 and 5, as well as from isolate No. 8, which contained a much lower level of apparent light and heavy chain association, were concentrated by ultrafiltration on a Centricon 30 filter (Amicon Corp.). The concentrated supernatants were run on a 7% polyacrylamide gel under non-reducing conditions, blotted to nitrocellulose, and probed with goat anti-human kappa antiserum followed by peroxidase-labeled rabbit anti-goat antiserum. The concentrated supernatants from isolates No. 1 and 5, but not from No. 8, contained a single immunoreactive band which comigrated with the purified chimeric L6 antibody from transfected Sp2/0 cells. These results suggested that isolates No. 1 and 5 were synthesizing and secreting assembled L6 chimeric antibody.

(5) Purification of chimeric L6 antibody from yeast culture supernatant.

[0238] In order to further characterize the H₂L₂-size protein secreted by the yeast and determine if this was assembled L6 chimeric antibody, a sufficient quantity of yeast-produced material was purified to allow the performance of various binding and functional assays. The pING1442 + 1441 transformant isolate No. 5 was grown for 58 hours at 30°C in a 10-liter fermentor using a synthetic medium (Table 8). The cells were initially grown in 9 liters of the column A medium until the glucose level fell below 1 g/L at which time they were fed with a total volume of 2.5 L of medium from column B. Glucose levels were maintained at 0.5 g/L during the remaining course of the fermentation. The cells were removed by centrifugation and the culture supernatant was analyzed by ELISA for the presence of light and heavy chain proteins and for association of the heavy and light chains. The supernatant contained approximately 250 µg/L of light chain, 240 µg/L of heavy chain, and 130 µg/L of heavy chain associated with light chain. The culture supernatants were next concentrated by ultrafiltration over a D.C. 10 unit (Amicon Corp.), filtered through 0.45 micron filter and concentrated over a YM30 filter (Amicon Corp.) to 250 ml. The concentrated supernatant was adjusted to pH 7.4 with KOH, brought

to 500 ml with PBS (10 mM sodium phosphate, pH 7.4, 150 mM sodium chloride) and loaded on a 1 ml protein A-Sepharose (Sigma) column, pre-equilibrated with PBS. The column was washed first with 20 ml PBS, followed by 10 ml 0.1 M sodium citrate, pH 3.5, then by 10 ml 0.1 M citric acid pH = 2.2. The pH 3.5 and 2.2 eluates were each collected in a tube containing 1 ml 2 M Tris base (Sigma). The bulk of the light and heavy chain immunoreactive proteins were in the pH 3.5 eluate which was next concentrated over a Centricon 30 (Amicon Corp.) to a final volume of 106 μ l. Analysis of this protein on non-reducing polyacrylamide gels using coomassie blue staining and immunoblotting with anti-human kappa antiserum (Sigma) to visualize the proteins revealed an H_2L_2 -size, 150 kilodaltons, protein band. This protein was purified away from other proteins by HPLC using an AB₈ 5-micron column equilibrated with buffer A (10 mM KPO₄, pH 6.8). After loading the sample on the column, the column was washed with buffer A for 10 minutes (flow rate = 1 ml/minute) and subjected to a linear gradient of 0% to 50% buffer B (250 mM KPO₄, pH 6.8) over 50 minutes at 1 ml/minute.

TABLE 8

MEDIUM USED FOR YEAST FERMENTATION TO PRODUCE SECRETED L6 CHIMERIC ANTIBODY ^a			
Ingredients		A ^b	B ^c
1.	Cerelose (Glucose)	119 g/l	538 g/l
2.	(NH ₄) ₂ SO ₄	13.9 g/l	83.3 g/l
3.	Thiamine HCL	0.011 g/l	0.05 g/l
4.	Biotin	0.00011 g/l	0.005 g/l
5.	Pantothenic acid	0.002 g/l	0.009 g/l
6.	Inositol	0.194 g/l	0.875 g/l
7.	H ₃ PO ₄	5.67 ml/l	25.5 ml/l
8.	KH ₂ PO ₄	5.78 g/l	26.0 g/l
9.	MgSO ₄ ·7H ₂ O	3.33 g/l	15.2 g/l
10.	CaCl ₂ ·2H ₂ O	0.33 g/l	1.5 g/l
11.	FeSO ₄ ·7H ₂ O	0.072 g/l	0.34 g/l
12.	ZnSO ₄ ·7H ₂ O	0.022 g/l	0.104 g/l
13.	MnCl ₂ ·4H ₂ O	0.0039 g/l	0.018 g/l
14.	CuSO ₄ ·5H ₂ O	0.0067 g/l	0.031 g/l
15.	Conc. H ₂ SO ₄	0.0056 ml/l	0.026 ml/l

a. Fermentation was performed as described in text.

b. Constituents of initial 9-liter batch.

c. Constituents of 2.5-liter feed batch.

[0239] The bulk of the protein resolved into a single large broad peak between 20 and 50 minutes as determined by absorbance at 280 nm. A second smaller peak was observed at 52-56 minutes, which corresponded to the normal elution position for chimeric L6 antibody from transfected Sp2/O cells. EUSA analysis of the column fractions revealed a major heavy + light chain cross-reactive peak corresponding to the U.V. absorbance peak at 52-56 minutes. Analysis of the 52-56 minute fractions on non-reducing SDS polyacrylamide gels using coomassie blue staining and immunoblotting revealed an essentially pure protein which co-migrated with L6 chimeric antibody purified from transfected Sp2/O cells.

(6) Studies performed on the chimeric L6 antibody secreted by yeast.

[0240] The purified yeast-derived antibody was assessed for function in several ways. First, the purified antibody was tested for its ability to bind directly to an L6 antigen-positive cell line. Second, the antibody was tested for its ability to inhibit binding of mouse L6 antibody to antigen-positive cells. Finally, the purified antibody was tested for two aspects of antibody function--the ability to mediate ADCC in the presence of human peripheral blood leukocytes and the ability to kill L6 positive tumor cells in the presence of human complement.

[0241] **Direct Binding Assay.** Cells from a human colon carcinoma line, 3347, which expresses approximately 5×10^5 molecules of the L6 antigen per cell on the cell surface, were used as targets. Cells from the T cell line, T51, were used as a negative control since they, according to previous testing, do not express detectable amounts of the L6 antigen. The target cells were first incubated for 30 min at 4°C with either the Sp2/O cell- or yeast-derived chimeric L6 antibody or with mouse L6 antibody standard purified from mouse ascites. This was followed by incubation with FITC-labeled goat-anti-human immunoglobulin for the chimeric antibodies or with FITC-labeled goat-anti-mouse immu-

noglobulin for the mouse standard. Both labeled antibodies were obtained from TAGO (Burlingame, CA) and used at a dilution of 1:50. Antibody binding to the cell surface was determined using a Coulter Model EPIC-C cell sorter.

[0244] As shown in Table 9, both the mammalian and yeast-derived chimeric L6 antibodies bound significantly, and to approximately the same extent, to the L6 positive 3347 line. They did not bind above background to the L6 negative TS1 line.

[0243] **Inhibition of Binding.** As the next step, the yeast chimeric L6 antibody and the Sp2/O cell-derived chimeric L6 antibody were tested for their ability to inhibit the binding of an FITC-labeled mouse L6 antibody to the surface of antigen-positive 3347 colon carcinoma cells.

[0244] Both the yeast-derived and Sp2/O-derived chimeric L6 antibodies inhibited the binding of labeled mouse L6 antibody and the binding curves were parallel. Based on the results of these studies, a rough estimate was made of antibody avidity. The avidity of the Sp2/O cell-derived chimeric L6 had been previously determined to be approximately 4×10^8 . The data indicated that there were no significant differences between the avidities of yeast-derived chimeric L6 and Sp2/O cell-derived chimeric L6 antibodies for the L6 antigen.

[0245] **Functional Assays.** A comparison was made between the ability of the yeast-derived chimeric L6, Sp2/O cell-derived chimeric L6 and standard mouse L6 antibodies to lyse L6 antigen-positive cells in the presence of human peripheral blood leukocytes as a source of effector cells mediating Antibody Dependent Cellular Cytotoxicity (ADCC). As shown in Table 10, the chimeric L6 from yeast was slightly better than Sp2/O-cell-derived chimeric L6 and as previously observed, both were superior to the standard mouse L6 in causing ADCC, as measured by a four-hour ^{51}Cr release test.

[0246] A comparison was next made between the yeast-derived chimeric L6, Sp2/O cell-derived chimeric L6 and standard mouse L6 antibodies for their abilities to lyse L6 antigen-positive cells by complement-dependent cytotoxicity (CDC) when human serum was used as the source of complement. The results of this comparison (Table 11) demonstrated that while both the Sp2/O-cell-derived chimeric L6 and standard mouse L6 antibodies exhibited high cytolytic activity, the yeast-derived L6 antibody failed to cause any cytotoxicity even at the highest antibody concentration. These results were unexpected and demonstrate that the yeast-derived antibody has new and unique properties.

(7) Conclusions

[0247] A process is disclosed by which yeast can be genetically engineered to secrete functional antibodies. The yeast-derived chimeric antibody in this example binds to the appropriate target antigen with approximately the same avidity as the chimeric antibody produced by lymphoid (Sp2/O) cells. The yeast-derived antibody also displays similar ADCC activity as does Sp2/O-derived antibody. Unlike the Sp2/O cell-derived antibody, the yeast-derived antibody displayed no CDC activity, thus demonstrating the new and unique properties of the yeast-derived antibody. This process should be applicable for the production of a variety of monoclonal antibodies and chimeric antibodies carrying chosen antigen binding domains linked to a chosen constant domain isotype. Genetically engineered antibodies and derivatives thereof produced in yeast also will exhibit novel functional properties, for example, the ability to selectively mediate target killing by ADCC without any detectable CDC activity. The technology described herein may also be suitable for the production of various other heterologous multimeric secreted proteins by genetically engineered yeast.

TABLE 9

BINDING ASSAYS OF CHIMERIC L6 ANTIBODY PRODUCED BY YEAST OR MOUSE Sp2/O CELLS ON AN L6 ANTIGEN-POSITIVE AND AN L6 ANTIGEN-NEGATIVE CELL LINE

Antibody ^a	Binding Ratio ^b for:	
	H3347 Cells (L6 +)	TS1 Cells (L6-)
Standard Mouse L6	95	1.0
Sp2/O Chimeric L6	116	1.0
Yeast Chimeric L6	116	1.0

a. All antibodies were used at a concentration of $10 \mu\text{g/ml}$.

b. The binding ratio is the number of times brighter a test sample is than a control sample treated with FITC-conjugated second antibody. Goat anti-mouse antibody was used as the second antibody for standard mouse L6 monoclonal antibody. Goat anti-human antibody was used as the second antibody for the yeast and Sp2/O chimeric L6 antibody.

TABLE 10

ADCC OF CHIMERIC L6 ANTIBODY DERIVED FROM YEAST OR Sp2/O CELLS AND STANDARD (MOUSE) L6 ANTIBODY ON COLON CARCINOMA CELL LINE 3347		
Antibody	Antibody Concentration ($\mu\text{g/ml}$)	% Cytolysis*
Standard mouse L6	5.0	42
	1.0	48
	1.0	96
	0.1	71
Sp2/O Chimeric L6	0.01	54
	0.001	37
	1.0	114
	0.1	108
Yeast Chimeric L6	0.01	76
	0.001	60
	0	23
	0	23

* The target cells had been labeled with ^{51}Cr and were exposed for four hours to a combination of MAbs and human peripheral blood leukocytes at 100 per target cell, and the release of ^{51}Cr was measured subsequently. The release of ^{51}Cr (after corrections of values for spontaneous release from untreated cells) is a measure of the percent cytotoxicity.

TABLE 11

HUMAN COMPLEMENT-DEPENDENT CYTOTOXIC EFFECTS OF CHIMERIC L6 ANTIBODY PRODUCED BY YEAST OR MOUSE Sp2/O CELLS ON COLON CARCINOMA CELL LINE 3347			
Antibody	Antibody Concentration ($\mu\text{g/ml}$)	Complement ^a (+ or -)	Percent Cytotoxicity
Standard mouse L6	5	+	122
	1	+	53
	5	-	1
Sp2/O Chimeric L6	5	+	73
	1	+	22
	0.1	+	5
	5	-	2
Yeast Chimeric L6	5	+	3
	1	+	2
	0.1	+	4
	5	-	2

^a Human serum from a healthy subject was used as the source of complement.

^b Complement-mediated cytotoxicity was measured by a four-hour ^{51}Cr -release assay.

EXAMPLE VI: Secretion of Functional Chimeric Fab from Yeast

[0248] The Fab portion of IgG consists of a single light chain molecule coupled by a disulfide bridge to a single truncated heavy chain molecule consisting of the variable region and C_{H1} (Figure 31). This heavy chain fragment is known as Fd. Fabs are potentially useful for a variety of therapeutic and diagnostic procedures. In addition, they are amenable to production by microbial fermentation.

[0249] The usual method for production of Fab involves the digestion of intact IgG with papain (see Figure 31) followed by purification of the Fab away from the Fc fragments generated in the digest. While this procedure is relatively straightforward and can result in high yields of Fab, it is somewhat time-consuming in that it first requires the production and purification of whole antibody followed by generation and, finally, purification of Fab. Furthermore, one-third of the whole antibody molecule—the Fc portion (Figure 31)—is not utilized.

[0250] The recent advances in gene cloning and site-specific mutagenesis technology make possible a more direct

and simple alternative approach for production of Fab molecules. In this approach, a stop codon is introduced in the heavy chain gene within the hinge region at approximately the codon for the amino acid at which papain digestion occurs. The Fab is then produced directly by simultaneous expression of both the light chain and Fd genes to produce their respective proteins which assemble and are secreted from the cell.

(1) Introduction of a stop codon in the hinge region of L6 chimeric heavy chain.

[0251] The strategy for introduction of a stop codon into the hinge region of L6 chimeric heavy chain is outlined in Figure 32A. The location of the stop codon within the hinge region and the DNA sequence of the mutagenesis primer are shown in Figure 32B. The stop codon placement corresponds to amino acid 226 in Figure 31. This procedure generated the plasmid pING1402 containing an Fd gene which codes for a protein consisting of 228 amino acids and extends six amino acids beyond the cysteine to which the light chain is coupled. The mutagenesis also introduced a unique BclI site at the stop codon which can be readily utilized for subsequent manipulations of the 3' end of Fd. These include, but are not necessarily limited to, removal of heavy chain 3' untranslated DNA as well as the engineering of various types of modifications of Fd including the addition of coding sequences for specific amino acids and the production of fusion proteins.

(2) Fusion of the mature Fd gene to yeast invertase signal sequence and shortened PGK promoter.

[0252] The strategy for fusion of the Fd gene to the yeast invertase signal sequence is outlined in Figure 33. This approach made use of the prior construction of the yeast invertase signal sequence-mature L6 heavy chain fusion (Figure 26) and utilized a unique ApaI site in the J region of the chimeric L6 heavy chain to replace the constant region in pING1415 consisting of C_H1 , C_H2 , and C_H3 with the constant region from pING1412 containing the stop codon in the hinge region. This procedure generated the plasmid, pING1418.

(3) Removal of non-yeast 3' untranslated DNA.

[0253] The introduction of a unique BclI site at the stop codon of the Fd chain provided a convenient method for removal of all non-yeast 3' untranslated DNA. This was accomplished using the strategy outlined in Figure 34, and generated the plasmid, pING1428.

[0254] Since the stop codon was introduced into the hinge region by site-specific mutagenesis of a heavy chain fragment cloned into M13, the possibility existed that unwanted mutations could have been introduced during the mutagenesis step. To ensure that such mutations were not present, an Fd gene fused to the Invertase signal sequence and shortened PGK promoter and consisting of known coding sequences was constructed using the strategy outlined in Figure 34, generating the plasmid, pING1444.

(4) Construction of yeast expression plasmids containing the chimeric L6 Fd gene from pING1444 fused to the PGK polyadenylation signal.

[0255] In order for yeast to produce an intact, functional Fab molecule, a balanced synthesis of both light and Fd-chain proteins must occur simultaneously within the cell. As described in Example V, one approach is to place the light chain and Fd genes on separate shuttle vectors containing separate selective markers and to transform these vectors into a yeast strain defective for both selective markers.

[0256] The Fd gene from pING1444 (Figure 34) was cloned as a BamHI-XhoI fragment into two medium copy number yeast-E. coli shuttle vectors containing sequences for replication in yeast and the PGK polyadenylation, transcription termination signal: pING804CVS for leu2 selection and pING1150 for ura3 selection (see Figures 29, 30). The two plasmids resulting from these constructions-pING1445 (ura3) and pING1446 (leu2) are shown in Figure 35.

(5) Secretion of chimeric L6 Fab from transformed yeast cells.

[0257] Two separate transformation experiments were performed in an attempt to obtain both light and Fd-chain synthesis in yeast cells. Four μ g each of pING1445 (Figure 35) and pING1441 (Figure 30) and separately of pING1446 (Figure 35) and pING1442 (Figure 30) were co-transformed into S. cerevisiae strain BB331c (MATa, ura3, leu2) by selecting for growth on SD agar (2% glucose, 0.67% yeast nitrogen base, 2% agar). Ura⁺ Leu⁺ transformants appeared at two to three days of incubation at 30°C.

[0258] Five colonies were inoculated from each plate into 6 ml SD broth supplemented with 50 mM sodium succinate, pH 5.5, and grown for 65 hours at 30°C. The cells were removed by centrifugation and analyzed by ELISA for the levels of light chain. The results of these assays revealed that the levels of light chain in the culture supernatants of

the pING1446 + pING1443 transformants were three to six times higher than the levels in the culture supernatants of the pING1445 + pING1441 transformants. The culture supernatants for each group of transformants were next concentrated by ultrafiltration on a Centricon 30 filter (Amicon Corp.) and run on a 10% polyacrylamide gel under non-reducing conditions. The proteins were blotted to nitrocellulose paper and probed with goat anti-human kappa antiserum followed by peroxidase-labeled rabbit-anti-goat antiserum. The concentrated supernatant from the pING1446 and pING1443 transformants contained a significant anti-kappa cross-reactive smear over a large portion of the blot with only a faint cross-reactive band at the position expected for the Fab protein. By comparison, the concentrated supernatants from pING1445 + pING1441 transformants contained relatively little smeared anti-human kappa cross-reactive protein on the blot. In addition, one of the five samples (No. 4) contained an especially intense, distinct anti-kappa cross-reactive band which migrated at the position expected for an Fab protein.

(6) Purification of chimeric L6 Fab from yeast culture supernatant.

[0259] To establish that the Fab-size anti-kappa cross-reactive protein secreted by the yeast is indeed L6 chimeric Fab protein required the purification of sufficient quantities for performance of binding assays. The pING1441 + pING1445 transformant isolate No. 4 was, therefore, grown in one liter of SD broth supplemented with 50 mM sodium succinate, pH 5.5, for 95 hours at 30°C. The cells were removed by centrifugation and the culture supernatant was analyzed by ELISA for the level of light chain protein. The supernatant contained approximately 130 µg/L of light chain protein. The culture supernatant was next concentrated by ultrafiltration over an Amicon YM30 filter to 20 ml. The concentrated supernatant was washed with 130 ml 10 mM potassium phosphate, pH 7.5 (buffer A) and re-concentrated over the YM30 filter to 12.5 ml. The concentrated supernatant was next brought to 54 ml with buffer A and loaded onto a 1.5 ml S-Sepharose column equilibrated with buffer A. The column was washed with 20 ml buffer A and subjected to a linear gradient of 0 to 200 mM sodium chloride in buffer A (40 ml total volume). ELISA analysis of the column fractions revealed a large anti-kappa cross-reactive peak between fractions 8 and 21 corresponding to a salt concentration of approximately 60 mM. These fractions were pooled, concentrated on Amicon YM10 and Centricon-10 filters (Amicon Corp.) to 51 µl and analyzed on non-reducing and reducing polyacrylamide gels using coomassie blue staining and Western blotting with anti-human kappa and anti-human Fab antisera. These analyses revealed an essentially pure protein which migrated at approximately 46 kd on the non-reducing gel and resolved into two bands running at approximately 23 and 24.5 kd on the reducing gel which corresponds to the predicted (based on amino acid sequence) molecular weights for light chain and Fd proteins, respectively. The smaller of the two bands strongly reacted with anti-human kappa antiserum on the Western blot. Both of the protein bands reacted with anti-human Fab antiserum on the Western blot.

(7) Studies performed on the chimeric L6 Fab secreted by yeast.

[0260] The primary activity of an Fab molecule is its ability to bind to the target antigen. The yeast-derived chimeric Fab was, therefore, tested for its ability to bind directly to an L6 antigen-positive cell line and for its ability to inhibit binding of mouse L6 antibody to antigen-positive cells.

[0261] Direct Binding Assay. Cells from the human colon carcinoma cell line 3347, which contains the L6 antigen at the cell surface, were used as targets. Cells from the antigen-negative cell line, T51, were used as a negative control. The target cells were first incubated for 30 minutes at 4°C with either yeast-derived chimeric L6 Fab, Sp2/O cell-derived chimeric L6 antibody, or with mouse L6 antibody. This was followed by incubation with FITC-labelled goat anti-human kappa immunoglobulin for the chimeric Fab, FITC-labelled goat anti-human IgG for chimeric antibody, or with FITC-labelled goat anti-mouse immunoglobulin for the mouse antibody. Both labelled antibodies were obtained from TAGO (Burlingame, CA) and used at a dilution of 1:50. Antibody binding to the cell surface was determined using a Coulter Model EPIC-C cell sorter.

[0262] As shown in Table 12, the yeast-derived chimeric L6 Fab bound to the L6 positive 3347 line. The yeast-derived chimeric L6 Fab did not bind above background to the L6 negative T51 line.

[0263] Inhibition of Binding. As the next step, we studied the extent to which graded doses of the yeast-derived chimeric L6 Fab or Sp2/O cell-derived chimeric L6 antibody could inhibit binding of an FITC-labelled mouse L6 antibody to the surface of antigen positive colon carcinoma 3347 cells.

[0264] The yeast-derived chimeric L6 Fab inhibited the binding of the directly labeled mouse L6 antibody. A higher concentration of the yeast L6 Fab, however, was required to achieve 50% inhibition of mouse L6 antibody binding to the target cells than was required for the same degree of binding inhibition by Sp2/O cell-derived chimeric L6 antibody.

(8) Conclusions

[0265] A process is disclosed by which yeast can be genetically engineered to secrete functional Fab domains of

immunoglobulins. The yeast-derived chimeric Fab in this example binds to the appropriate target antigen. Such Fab molecules provide convenient targeting agents for a variety of diagnostic and therapeutic uses. This process also demonstrates the feasibility of secretion of heterologous heterodimeric molecules from yeast.

TABLE 12

BINDING ASSAYS OF CHIMERIC L6 FAB PRODUCED BY YEAST ON AN L6 ANTIGEN-POSITIVE AND AN L6 ANTIGEN-NEGATIVE CELL LINE

Antibody ^a	Binding Ratio ^b for:	
	3347 Cells (L6 +)	T51 Cells (L6-)
Sp2/O Chimeric L6	103	1
Yeast Chimeric L6 Fab	32	1

a. All antibodies were used at a concentration of 10 µg/ml.

b. The binding ratio is the number of times brighter a test sample is than a control sample treated with FITC-conjugated second antibody. Goat anti-human antibody was used as the second antibody for the Sp2/O chimeric L6 antibody and goat-anti-human kappa antibody was used as the second antibody for the yeast Fab.

EXAMPLE VII: Secretion of Functional Chimeric Fab Molecules From Bacteria

[0266] Bacteria are suited for production of chimeric antibodies expressed from mammalian cDNA since entire coding sequences can be expressed from well characterized promoters. *Escherichia coli* is one of many useful bacterial species for production of foreign proteins (Holland, I.B., et al., *BioTechnology* 4:427 (1986)), since a wealth of genetic information is available for optimization of its gene expression. *E. coli* can be used for production of foreign proteins internally or for secretion of proteins out of the cytoplasm, where they most often accumulate in the periplasmic space (Gray et al., *Gene* 39:247 (1985); Oka et al., *Proc. Natl. Acad. Sci. USA* 82:7212 (1985)). Secretion from the *E. coli* cytoplasm has been observed for many proteins and requires a signal sequence. Proteins produced internally in bacteria are often not folded properly and precipitate into subcellular particles called inclusion bodies (Schoener et al., *BioTechnology* 3:151 (1985)). Protein secreted from bacteria, however, is often folded properly and assumes native secondary and tertiary structures (Hsiung et al., *BioTechnology* 4:991 (1986)). Although immunoglobulin peptides have been synthesized in genetically engineered *E. coli* (Cabilly et al., *Proc. Natl. Acad. Sci. USA* 81:3273 (1984); Liu et al., *Proc. Natl. Acad. Sci. USA* 81:5369 (1984); Boss et al., *Nucl. Acids Res.* 12:3791 (1984)), there are no reports of secretion of these peptides from *E. coli* as functional antibodies or antibody fragments.

[0267] An Fab molecule consists of two nonidentical protein chains linked by a single disulfide bridge. These two chains are the intact antibody light chain and the V, J, and C_{H1} portions of the antibody heavy chain, Fd. The proper cDNA clones for the L6 chimeric light and Fd gene have already been identified. In this example, these cDNA clones were organized into a single bacterial operon (a dicistronic message) as gene fusions to the pectate lyase (*pelB*) gene leader sequence from *Erwinia carotovora* (Lei et al., *J. Bacteriol.*, in press (1987)) and expressed from either of two strong, regulated promoters. The result is a system for the simultaneous expression of two protein chains in *E. coli*, and the secretion of immunologically active, properly assembled Fab of L6 chimeric antibody into the culture growth media.

A. Construction of *E. coli* expression systems for L6 Chimeric Fab.

1. Assembly of the *pelB* leader sequence cassette.

[0268] *Erwinia carotovora* EC codes for several pectate lyases (polygalacturonic acid trans-eliminase) (Lei et al., *Gene* 35:63 (1985)). Three pectate lyase genes have been cloned, and the DNA sequence of these genes has been determined. When cloned into *E. coli* under the control of a strong promoter, the *pelB* gene is expressed and large quantities of pectate lyase accumulate in the periplasmic space. The *pelB* signal sequence functions efficiently in *E. coli* and was used as a secretion signal for antibody genes in this example. The nucleotide sequence surrounding the signal sequence of the *pelB* gene is shown in Figure 36a.

[0269] The *pelB* signal sequence contains a *HaeIII* restriction site at amino acid 22, adjacent to the signal peptidase cleavage site: ala-ala. Plasmid pSS1004 (Lei et al., *J. Bacteriol.*, in press (1987)), containing the *pelB* gene in plasmid vector pUC3 (Vieira and Messing, *Gene* 19:259 (1982)), was digested with *HaeIII* and *EcoRI*. This DNA was ligated with an eight base pair *SstI* linker to *SspI* and *EcoRI* cut pBR322. The resulting plasmid contained a 300 bp fragment which included the 22 amino acid leader sequence of *pelB* and about 230 bp of upstream *E. carotovora* DNA. This plasmid pING173, contains an insert that upon digestion with *SstI* and treatment with T4 DNA polymerase can be

ligated directly to a DNA fragment flanked by the first amino acid of a mature coding sequence for any gene to generate a protein fusion containing a functional bacterial leader sequence in frame with the incoming gene. The SstI to EcoRI restriction fragment in pING173 was cloned into pUC18 (Yanich-Perron *et al.*, Gene 33: 103 (1985)) to generate pRR175, which contains the pelB leader and adjacent upstream non-coding sequence (including a ribosome binding site) downstream of the lac promoter. The construction of pRR175 is outlined in Figure 36b.

2. Preparation of chimeric L6 light gene for bacterial expression.

[0270] The intact L6 chimeric light chain gene containing an AatII restriction site at the signal sequence processing site and a unique BglII site downstream of the gene was excised from the yeast expression plasmid pING1298 (Figure 25a) as a 1200 bp DNA fragment. This fragment was inserted into plasmid pRR175. The resulting plasmid, pRR177-8, contained an in-frame fusion of the pelB leader and the L6 light chain downstream of the lac promoter residing in the parent plasmid. A number of derivatives of this plasmid were constructed to delete noncoding sequences from both the 5' and 3' ends of the pelB:light chain gene fusion in pRR177-8. Upstream noncoding sequences were deleted making use of an NdeI restriction site at -48 bp from the pelB leader sequence initiation codon (Figure 36) generating pRR180-2. The 3' noncoding sequences were eliminated by substituting a fragment from the plasmid optimized for L6 light chain expression in yeast, pING1431 (see Figure 27a), into pRR179 to generate pRR191. Another plasmid, pRR190, is similar to pRR191 but contains 90 bp of noncoding eukaryotic DNA at the 3' end of the light chain gene. These constructions are shown in Figure 37.

3. Preparation of chimeric L6 Fd gene for bacterial expression.

[0271] The intact L6 chimeric Fd gene containing an SstI restriction site at the signal sequence processing site, a BclI site introduced by site directed mutagenesis (Figure 32a, b) and creating a termination codon at amino acid 226, and a unique BamHI restriction site downstream of the gene was excised from the plasmid pING1406 (Figure 33) as a 860 bp DNA fragment. This DNA fragment was inserted into plasmid pRR175 generating an in-frame fusion of the pelB leader sequence and the L6 Fd gene downstream of the lac promoter, pRR178-5. A number of derivatives were constructed to delete noncoding sequences from both the 5' and 3' ends of the sequence contained in pRR178-5. The 3' noncoding sequences were eliminated by substituting a restriction fragment from the plasmid optimized for L6 Fd expression in yeast, pING1428 (Figure 34), which contains an XhoI linker immediately following the termination codon of the Fd gene, generating plasmid pRR186. Removal of E. carotovora DNA sequences upstream of the NdeI site at -48 from the leader sequence generated plasmid pRR196. The construction of these plasmids is shown in Figure 38.

4. Multicistronic expression system for light chain and Fd gene.

[0272] For production of bacterially derived Fab, both light chain and Fd need to be produced simultaneously within the cell. Using the plasmids constructed with each of these genes separately, a series of expression vectors were constructed that contain both genes aligned so that transcription from a single promoter will specify both genes. This was done in a way that minimized the noncoding DNA between the two genes to 60 bp. Each gene has a ribosome binding site needed for translation initiation and the identical DNA sequence from -48 to the pelB leader: antibody gene junction. Several cloning steps were required to align the two genes together. A portion of the light chain gene linked to the pelB leader in pRR180-2 was cloned downstream of the Fd gene in pRR186 to generate pFK100. The remainder of the light chain gene was subcloned into pFK100 from pRR177-8 to generate pFK101. Similarly, DNA fragments containing 3' deletions of eukaryotic sequences from pRR190 and pRR191 were cloned into pFK101 generating pFK103 and pFK102 respectively. DNA fragments from pRR192 and pFK101 were ligated to generate pFK104 which contains a deletion of sequences upstream of -48 bp from the Fd gene. Maps of the Fd and light chain gene cassettes in these plasmids are shown in Figure 39.

5. Placement of the dicistronic message for light chain and Fd under the control of inducible promoters.

[0273] Plasmids pFK101, pFK102, pFK103, and pFK104 contain Fd and light chain genes cloned sequentially under the control of the lac promoter in vector pUC18 or pUC19. In E. coli strains such as JM103 F^{+lac^Q (Messing *et al.*, Nucl. Acids. Res. 9:309 (1981)), the amount of light chain that accumulates in the periplasm is not affected by the lac promoter inducing agent isopropyl β-D-thiogalactopyranoside (IPTG), see Table 13. In addition, bacterial growth is slower (compared to cells containing pUC18), and bacterial colonies exhibit an altered morphology being small, dry and rough, suggesting that constitutive foreign gene expression is deleterious to cell growth. Two strategies were used to place this gene cassette under more tightly regulated promoters.}

[0274] First, a PstI to EcoRI fragment from pFK104 was ligated to pIT206 to place the Fd and light chain gene cassette

under the direct control of the *Salmonella typhimurium* *araB* promoter, a well characterized, strong promoter in *E. coli*. A restriction map of pIT206 and construction of pIT104 is shown in Figure 40. Use of the *araB* promoter and its regulatory protein *araC* for the expression of bacterial genes is described in U.S. Patent Applications 695,309 filed January 28, 1985, and 797,472, filed November 13, 1985. As is seen in Table 14, the resulting plasmid, pIT104, is now regulated for the synthesis of light chain by the addition of arabinose to the culture growth media. At least 10 fold induction is effected by arabinose addition. Although Fab secreted into the growth medium increases more than 10 fold, cell growth stops after induction with arabinose. This confirms that high level expression of the Fab genes is deleterious to cell growth. Bacterial colonies harboring pIT104 are phenotypically indistinguishable from *E. coli* harboring pIT206 when grown in the absence of arabinose.

[0275] Second, a DNA fragment containing the *lacI* gene, a repressor of the *lac* promoter, was cloned into the high copy expression vector pFK102. Expression of *lacI* from a high copy number vector is useful to regulate expression of the *lac* promoter on a high copy number vector (Russell et al., Plasmid, in press (1987); Hsuing et al., *Biotechnology* 4:991 (1986)). A 1.7 kb *EcoRI* fragment containing the *lacI* gene on pMC9 (Calos et al., *Proc. Natl. Acad. Sci. USA* 80:3015 (1983)) was excised, filled in with T4 polymerase to blunt ends, ligated with *PstI* linkers and cloned into the unique *PstI* site of pFK102 to generate pFK102*lacI*. The map of pFK102*lacI* is shown in Figure 40b. The selection procedure used to identify the correct clone assured that the resulting plasmid, pFK102*lacI*, contained a functionally repressed *lac* promoter. All white or light pink colonies on MacConkey-lactose plates contained plasmids with *lacI* inserts while transformants containing pFK102 alone were red, indicating the functional repression of the *lac* promoter by the high copy number *lacI* gene. Table 14 shows that expression of bacterial Fab from cells containing pFK102*lacI* is similar to expression from pFK102. Unlike cells containing pFK102, which formed aberrant colonies and grew slowly in broth culture, cells containing pFK102*lacI* resembled those containing pUC18.

B. Expression, SDS-PAGE, and Purification of Bacterially Produced Fab

1. Growth of *E. coli* harboring cloned antibody genes.

[0276] Plasmid DNA was transformed into either *E. coli* JM103 or MC1061 by standard *E. coli* transformation procedures. Bacterial cultures were grown in TYE (tryptone 1.5%, yeast extract 1.0%, and NaCl 0.5%) supplemented with the appropriate antibiotics (penicillin 250 ug/ml, tetracycline 15 ug/ml). Bacterial cultures were grown in volumes of 5 ml to 1 liter at 37°C to an optical density OD₆₀₀ = 0.8 (approximately 4 X 10⁸ cell/ml) and aliquots were induced with IPTG (0.2 mM), lactose (1.0%), or arabinose (1.0%). Cultures were grown for an additional time period of 4 to 21 hr. Portions of each culture were analyzed for light chain production. Protein was released from the periplasmic space of *E. coli* cells by osmotic shock as described (Yanagida et al., *J. Bacteriol.* 166:937 (1986)). Alternatively, culture supernatants were assayed directly for the presence of antibody chains.

[0277] Quantitation of L6 light chain was by ELISA with goat anti-human Kappa light chain antibody (Cappel, Malvern, PA). Fd could be detected by ELISA with mouse monoclonal anti-human Fd antibody (Calbiochem, San Diego, CA). Table 13 shows representative data for expression of light chain reactive material in *E. coli* periplasmic extracts. Light chain is secreted from the bacterial cytoplasm into the periplasm. Antibody chains are also released from the bacteria into the culture supernatant. This is an unusual discovery and may be a unique property of the L6 Fab among eukaryotic proteins expressed in *E. coli*. Under certain conditions, however, bacterial proteins are known to be released from *E. coli* (Abrahmsen et al., *Nucl. Acids Res.* 14:7487 (1986); Pages et al., *J. Bacteriol.* 169:1386 (1986)). Table 14 compares the amount of light chain secreted into the periplasm with the amount secreted into the culture supernatant. Light chain reactive material is present in plasmid containing cultures harboring cloned light chain alone or light chain plus Fd. The best producers of Fab (pFK102, pFK104, and pFK102*lacI*) typically secrete 300 - 1000 ng/ml of ELISA reactive light chain into the culture media. A separate construct was made in which the light chain gene is followed by the Fd gene (pFK107). This construct directs synthesis and secretion of Fab at similar levels to the constructs with the genes in the inverse order. Thus, the gene order is not critical for secretion of Fab.

2. SDS-PAGE of bacterially produced chimeric L6 light chain and Fd.

[0278] Bacterially produced antibody chains were analyzed by polyacrylamide gel electrophoresis under reducing and non-reducing conditions. Protein extracts of lysed whole bacterial cells, protein released from the periplasmic space by osmotic shock, and protein secreted into the culture supernatant were analyzed electrophoretically. Transfer of gel separated protein under full reducing conditions to nitrocellulose and immunological staining with goat anti-human light chain antibody by Western analysis revealed that a protein of the same molecular weight as authentic L6 chimeric light chain was present (about 23 Kd). Analysis of protein samples by SDS-PAGE under non-reducing conditions showed that extracts from cells harboring a plasmid with the light chain gene alone (pRR191 or pRR190) contained a large proportion of the light chain reactive material associated into a higher molecular weight form. Much of

this material ran at about 46 Kd in what is likely to be a light chain dimer. Light chain dimers have been observed from myeloma cells producing only light chain. There are also other immunoreactive protein bands that may represent non-specific disulfide formation between light chain and *E. coli* proteins. Protein samples (periplasmic extracts or culture supernatants) from *E. coli* cells harboring both the light chain and the Fd genes contain a light chain reactive band at about 48 Kd when separated under non-reducing gel conditions which runs at a slightly higher molecular weight than the bacterial light chain dimer. This material is bacterially produced L6 chimeric Fab. In *E. coli* harboring pFK102lacI, pFK101, pFK102, pFK103, or pFK104 the 48 Kd band observed on an SDS gel run under non-reducing conditions is the most prominent immunoreactive specie. In addition, the background smear of immunoreactive proteins seen in extracts containing the light chain only is greatly reduced in extracts from cells containing both light chain and Fd.

3. Purification of bacterially produced chimeric L6 Fab.

[0279] Immunologically and functionally active (see below) bacterial Fab was purified from either culture supernatants or periplasmic protein extracts of *E. coli* harboring pFK102lacI or pT104. For purification of periplasmic material, the periplasmic fraction from 1 liter of cells induced for 4 hours was released into 50 ml of distilled water. This material was centrifuged for 20 minutes at 5000 g and filtered through a 0.45 µm filter. The periplasmic extract was then concentrated over a YM10 membrane (Amicon) to about 5 ml. This material was diluted 8 fold into starting buffer (10 mM K₂HPO₄, pH 7.5) and applied to a 1 ml S-Sepharose column at a flow rate of 1.0 ml/min. The column was washed with 25 ml of starting buffer and eluted with a 0 to 200 mM NaCl gradient in starting buffer (200 ml total volume). The immunoreactive gradient peak was pooled (elution was at about 100mM) and concentrated on a Centricon 10. Purified Fab was stored in PBS + 2.0% BSA.

[0280] For purification of secreted Fab from 1 liter of bacterial culture supernatant, the cells were removed by centrifugation after growth for 21 hours with inducing agents and the supernatant was filtered through a 0.45 µm filter. The media was concentrated over a YM10 membrane (Amicon) to about 16 ml, then diluted with 10 mM K₂HPO₄ to 105 ml. This material was applied to a 1.6 ml S-Sepharose column and eluted with a 0 to 200mM NaCl gradient in 40 ml. Fab recovered from S-Sepharose chromatography was greater than 70% pure as determined by densitometry tracing of a nonreducing, coomassie stained, 10% acrylamide gel. The Fab purified from bacterial culture supernatants resolves into two major protein bands of about 23 Kd and 24.5 Kd on a 15% reducing gel. The molecular weight of Fd and light chain based on the DNA sequence are 24.5 Kd and 23 Kd which corresponds well to the observed protein sizes. The smaller of the two bands strongly reacted with goat anti-human Kappa light chain antiserum on a Western blot. Bacterial Fab purified from either the periplasmic space or bacterial culture supernatants are indistinguishable by all analytical criteria tested here.

4. Functional binding activity of bacterially produced chimeric L6 Fab to the L6 antigen.

[0281] Bacterially produced Fab purified by S-Sepharose chromatography was tested for binding to L6 antigen containing cells. As shown in Table 15, bacterial Fab binds specifically to the human colon carcinoma cell line 3347. Cells from the T cell line T51 were used as a negative control. Target cells were incubated for 30 minutes at 4°C with bacterially produced L6 chimeric Fab, intact L6 chimeric antibody produced in Sp2/0 cells, or mouse L6 antibody purified from mouse ascites. This was followed by incubation with FITC-labelled goat anti-human light chain antibody for Fab detection, FITC-labelled goat anti-human immunoglobulin for chimeric antibody detection, or with FITC-labelled goat antimurine immunoglobulin for mouse antibody detection. Antibody binding to the cell surface was determined using a Coulter Model EPIC-C cell sorter.

[0282] Bacterially produced Fab also exhibits characteristic binding inhibition of FITC-labelled mouse L6 antibody to the surface of antigen positive 3347 colon carcinoma cells. Bacterially produced Fab and Sp2/0 derived chimeric L6 have similar binding inhibition profiles, thereby suggesting that the avidity of bacterially produced Fab and Sp2/0 derived chimeric L6 are similar.

Conclusions

[0283] A process is disclosed whereby *E. coli* has been used as a host to produce functionally active Fab domains of immunoglobulins and to secrete these into the periplasmic space and also in the culture medium. This molecule exhibits binding properties expected of a properly assembled antibody recognition site. This technology can be used to express antibody genes with other binding specificities in *E. coli*.

1. Proteins encoded by modified antibody cDNA clones can be secreted from bacteria using a signal sequence
2. Two antibody genes can be expressed from a single bacterial promoter as a dicistronic message.
3. Two foreign proteins (in this example antibody light chain and Fd) can assemble properly, i.e., assume correct

secondary, tertiary, and quaternary structure when secreted from bacteria.

4. At least two, and probably many bacterial promoters can be used for expression of antibody genes.

5. This example is a general method whereby genes encoding other antibody chains can be expressed together as a dicistronic message; these include either light chain and Fd genes or light chain and intact heavy chain genes.

6. The gene order with respect to the promoter is not important in the ability of E. coli to produce Fab. A construct of the Fd gene followed by the light chain works as well as the genes organized in the inverse order.

7. Fab can be secreted from E. coli into the culture supernatant where it is stable and can be purified. Most Fab chains that pass the cytoplasmic membrane are secreted into the culture supernatant.

Microorganism Deposits

[0284] Saccharomyces cerevisiae BB331C (41/42-5), G187 was deposited at the ATCC on July 9, 1987 and given access number 20856. Escherichia coli JM 103 (pFK1021 aci), G186 was also deposited therein on the same date and given access number 67457. Both deposits were under the Budapest Treaty.

TABLE 13
QUANTITATION OF LIGHT CHAIN FROM E. COLI PERIPLASM

plasmid	ng/ml of culture		plasmid	ng/ml of culture	
	-	+		-	+
pRR175	0	0	pFK101	38	28
pRR177-8	8.5	11	pFK102	68	55
pRR180	388	412	pFK103	38	45
pRR190	200	241	pFK104	81	88
pRR191	483	772			

E. coli JM103 or MC1061 (results similar) was transformed with each plasmid. Fresh transformants were cultured in TYE at 37°C to an OD₆₀₀ = 0.8. Cultures were divided and the inducer (IPTG) was added to 0.2 mM to one aliquot (- or + IPTG). Cells were grown at 37°C for 4 hours. Periplasmic protein extracts were prepared, and aliquots were tested for light chain by ELISA with goat anti human kappa antibody. Each value is the average of at least two separate experiments. Removal of non-coding sequences both 5' and 3' to the antibody gene effected in increase on light chain accumulation in the periplasm.

TABLE 14

ACCUMULATION OF LIGHT CHAIN IN THE SUPERNATANT AND PERIPLASM AFTER INDUCTION						
plasmid	Inducer	Supernatant			Periplasm	
		4 hr	21 hr	4 hr	21 hr	
pRR180	-	0	nd	200	nd	nd
pRK180	+	5	188	241	nd	nd
pFK102	-	12	nd	89	nd	nd
pFK104	+	57	829	55	40	40
pFK104	-	13	nd	91	nd	nd
pFK102lac	+	150	290	66	35	35
pFK102lac	-	25	360	50	100	100
pT104	+	72	608	37	40	40
pT104	-	13	nd	10	nd	nd
pT104	+	150	216	18	35	35

Plasmid containing *E. coli* strains were grown, prepared, and assayed as described in Table 13. For pRR180, pFK102, pFK104, and pFK102lac cells were induced with 0.2 mM IPTG; pT104 was induced with 1% arabinose. Each value is the average of at least two separate experiments. For analyses of *E. coli* culture supernatants, bacteria were removed by centrifugation and culture supernatants were passed through a 0.45 μ m filter. Values are expressed in ng/ml of culture.

nd - not determined

TABLE 15
BINDING ASSAYS OF BACTERIAL Fab

Antibody	Binding ratio*	
	3347 cells L6 +	T51 cells L6-
Standard mouse L6	95	1
Sp2/0 chimeric L6	116	1
Bacterial L6 Fab	54	1
Standard L6 Fab	16	1
Standard L6 Fab was prepared by enzymatic digestion of mouse L6 antibody.		

* The binding ratio is the number of times brighter a test sample is than a control sample treated with FITC-conjugated second antibody.

Claims

1. A polynucleotide molecule comprising a prokaryotic promoter region in operable linkage to a dicistronic transcription unit, the unit encoding a heavy chain immunoglobulin, or a fragment thereof, and a light chain immunoglobulin, the heavy and light chains being separately operably linked to a sequence coding for a polypeptide secretion signal.
2. A molecule according to Claim 1 wherein the heavy and light chains are chimeric.
3. A vector comprising recombinant DNA and a polynucleotide molecule as defined in any of Claim 1 or Claim 2 and capable of expressing the polynucleotide molecule in a prokaryotic host.
4. A prokaryotic host capable of producing an immunoglobulin transfected with a vector as defined in Claim 3.

Patentansprüche

1. Polynucleotidmolekül, welches eine prokaryotische Promotorregion in funktioneller Verknüpfung mit einer dicistronischen Transkriptionseinheit aufweist, wobei die Einheit eine schwere Kette eines Immunglobulins oder ein Fragment davon und eine leichte Kette eines Immunglobulins codiert, wobei die schweren und leichten Ketten getrennt funktionell mit einer Sequenz verknüpft sind, die Polypeptidsekretionssignal codiert.
2. Molekül nach Anspruch 1, bei welchem die schweren und die leichten Ketten chimär sind.
3. Vektor, welcher rekombinante DNA und ein Polynucleotidmolekül gemäß der Definition in einem der Ansprüche 1 oder 2 enthält, und welcher fähig ist, das Polynucleotidmolekül in einem prokaryotischen Wirt zur Expression zu bringen.
4. Prokaryotischer Wirt, welcher in der Lage ist, ein Immunglobulin zu erzeugen, welcher mit einem Vektor gemäß der Definition in Anspruch 3 transfiziert ist.

Revendications

1. Molécule polynucléotidique comprenant une région promoteur provenant d'un procaryote, en liaison opérationnelle avec une unité de transcription dicistronique, l'unité codant pour une immunoglobuline à chaîne lourde ou un fragment de celle-ci, et une immunoglobuline à chaîne légère, les chaînes lourde et légère étant liées séparément de manière opérationnelle à une séquence codant pour un signal de sécrétion de polypeptide.
2. Molécule suivant la revendication 1, dans laquelle les chaînes lourde et légère sont chimériques.
3. Vecteur comprenant un ADN recombinant et une molécule polynucléotidique comme défini dans l'une quelconque des revendications 1 et 2, et capable d'exprimer la molécule polynucléotidique dans un hôte procaryote.

4. Hôte procaryote capable de produire une immunoglobuline, transfecté par un vecteur défini dans la revendication 3.

5

10

15

20

25

30

35

40

45

50

55




FIG. 1

Ig heavy chain J-C regionhuman heavy chain J regions

J1 CH1

JH1 CCTGAATACTTCCAGCACTGGGGCCAGGGCACCCCTGGTCACCGTCTCCTCAG
 JH2 CTACTGGTACTTCGATCTCTGGGGCCGTGGGCACCCTGGTCACTGTCTCCTCAG
 JH3 ATGCTTTTGATGTCTGGGGCCAAAGGACAAATGGTCAACGTCCTCTCAG
 JH4 ACTACTTTTGACTACTGGGGCCAAAGGAACCTGGTCAACGTCCTCCTCAG
 JH5 ACACCTGGTTCGACTCCTGGGGCCAAAGGAACCTGGTCAACGTCCTCCTCAG
 JH6 AT(TAC)₅GGTATGACGCTCTGGGGCCAAAGGACCAACGTCACCGTCTCCTCAG
Consensus TCGACCTCTGGGGCCAAAGGAACCTGGTCAACGTCCTCCTCAG

mouse heavy chain J regions

J1 CH1

JH1 TACTGGTACTTCGATGTCTGGGGCCAGGGACCAACCGTCAACGTCCTCCTCAG
 JH2 TACTTTGACTACTGGGGCCAAAGGCACCCTCTCACAGTCTCCTCAG
 JH3 CCTGGTTTGCTTACTGGGGCCAAAGGACTCTGGTCACTGTCTCTGCAG
 JH4 TACTATGCTATGGACTACTGGGGTCAAGGAACCTCAGTCAACGTCCTCCTCAG
Consensus TTTGACTACTGGGGCCAAAGGACCAACCGTCAACGTCCTCCTCAG

Ig light chain J-C regionhuman Kappa J region

J1C

JK1 GGACGTTCCGCCAAGGGACCAAGGTGGAATCAAAC
 JK2 ACACCTTTTGGCCAGGGACCAAGCTGGAGATCAAAC
 JK3 TCACCTTTCCGCCCTGGGACCAAGGTGAGATCAAAC
 JK4 TCACCTTTCCGCCGAGGGACCAAGGTGAGATCAAAC
 JK5 TCACCTTTCCGCCAAGGGACCAAGCTGGAGATCAAAC
Consensus TTCGCCCAAGGGACCAAGGTGAGATCAAAC

mouse Kappa J region

J1C

JK1 TCGACGTTCCGTGGAGGCACCAAGCTGGAATCAAAC
 JK2 TACACGTTCCGAGGGGGACCAAGCTGGAATCAAAC
 JK3 TTCACATTCAGTGATGGGACCAAGCTGGAATCAAAC
 JK4 TTCACGTTCCGCTCGGGGACCAAGTTGGAATCAAAC
 JK5 CTCACGTTCCGTGCTGGGACCAAGCTGGAGCTGAAAC
Consensus TTCGTTGGGGGACCAAGCTGGAATCAAAC
 UIG[MJK] 3' TGGTTCGACCTTTATTTTG 5'

human Lambda pseudo J region

J1C

JPSL1 CACATGTTTGGCAGCAAGACCCAGCCCACTGTCTTAG

mouse Lambda J region

J1C

JL1 TGGGTGTTCCGTGGAGGAACCAACTGACTGTCTCTAG
 JL2 TATGTTTTCCGGCGGTGGAACCAAGGTCACTGTCTCTAG
 JL3 TTTATTTTCCGCGAGTGAACCAAGGTCACTGTCTCTAG
Consensus TTCGGCGGTGGAACCAAGGTCACTGTCTCTAG

FIG. 2




FIG. 3





FIG. 5

A. SYNTHESIS OF HUMAN IgG1 GENES
a HUMAN IgG1 HEAVY CHAIN STRUCTURE



B. A HUMAN IgG1 CONSTANT REGION CLONING VECTOR FOR V REGION MODULE INSERTION





FIG. 4

A. SYNTHESIS OF HUMAN IgK GENES

a. HUMAN IgK LIGHT CHAIN STRUCTURE



b. cDNA CLONES



B. CONSTRUCTION OF A HUMAN C_κ REGION CLONING VECTOR

FIG. 6

Primers Designed for Ig V Region Synthesis

A. Ig Heavy Chain J-C Region

Human IgG1 pGMH-6
 — J Region — | — IgG1 CH1 Region —
 GGTCACCGTCTCTCTCAG CCTCCACCAAGGGCCCATC
Bs/EI

Mouse Heavy Chain J Regions and Primers

	N	Mismatches			JH4
		JH1	JH2	JH3	
JH1 TACTGGTACTTCGATGCTCTGGGCGCAGGGACACCGGTACCGTCTCTCTCAG [JH1] GCCATGGCAGAGGATCGGT	21	0	4	4	1
JH2 TACTTTGACTACTGGGCGCAAGGCACCACTCTCACAGTCTCTCTCAG [JH2] GAGAGTGTACAGCAGTCGGT	21	4	0	7	4
JH3 CCTGGTTTGCTTACTGGGCGCAAGGGACTCTGTCTCTCTCTCTCGAG [JH3] ACCACTGACACAGACGTCGGT TCCCTGAGAGCCAGTGGCAGAG ACCACTGGCAGAG <u>Bs/EI</u>	21 21 13	4 3 1	7 7 4	0 1 1	5 5 2
JH4 TACTATGCTATGGACTACTGGGTCAAGGACCTCAGTCACCGTCTCTCTCAG [JH4] GTCATGGCAGAGGATCGGT	21	1	4	5	0

FIG. 7A

Primers Designed for Ig V Region Synthesis

B. Ig Kappa Chain J-C Region

-J Region-----IgK Constant Region-----
 CTGGAGATCAAC GAACGTGGCTGCACCATCTGCTTCATCTTCC
TCAGCAAC GAACGTGGCTGCACCATCTGCTTCATCTTCC
Bcl

Human kappa pK1-3
 PING2001

House Kappa J Regions and Primers

	N	JK1	JK2	JK4	JK5	Mismatches
JK1 TGGAGTTCGGTGGAGGACCAAGCTGGGAATCAAC {5JK1} GCAGCCACCTCCGTGG	17	0	3	6	3	
JK2 TACAGTTCGGAGGGGGACCAAGCTGGGAATCAAC {JK2BGL1} CCCTGGTTCGACCTCTAGATT <i>Bgl</i>	21	3	3	5	3	
JK4 ATCAGTTCGGCTCGGGACAAAAGTTGGGAATCAAC {5JK4} GCAGCCGAGCCCTGT {JK4BGL1} GCCCTGTTTCAACCTCTAGATT <i>Bgl</i>	17	6	4	0	4	
	23	7	6	3	6	
JK5 CTCAGTTCGGTCTGGGACCAAGCTGGGCTGAAC {5JK5} GCAGCCACGACCTGG	17	3	3	4	0	

FIG7A(cont.)

C. Mouse Variable Region Consensus Primers

mouse heavy chain J segments

JH1 T A C T G G T A C T T C G A T G T C T G G G C C A G G G A C C A C G G T C C T C A
 JH2 T A C T T T G G C T A C T G G G C C A G G G A C C A C G G T C C T C A
 JH3 C C T G G T T T G C T A C T G G G C C A G G G A C C A C G G T C T G C A
 JH4 T A C T A T G C T A T G G A C T A C T G G G T C A G G G A C C A C G G T C C T C A
 consensus primer: UIG-H A G G A C C A C G G T C A C C G T C T C

BseEII

T C C C T G G T G C C A G T G G C A G A G
 3' 5'

mouse light chain J segments

JK1 T G G A C G T T C G G T G G A G G A C C A A G C T C G A A T C A A A
 JK2 T A C A C G T T C G G A G G G G G A C C A A G C T C G A A T A A A A
 JK4 T T C A C G T T C G G C T C G G G G A C C A A G C T C G A A T A A A A
 JK5 T T C A C G T T C G G T G C T G G G A C C A A G C T C G A C T G A A A
 consensus primer: UIG-K G G G A C C A A G C T T G A G

HindIII

C C C T G G T T C G A A C T C
 3' 5'
 G G A G G G A C C A A G G T G G A T G A A
 C - T - - - - -
 HindIII

pCM.60

D. Mouse y2a J/C Junction Primer

MJ_H2-Apa1 T G T C A G A G G A G T C G G T C G T G T T C C C G G G T A
 3' 5'
 Apa1

FIG. 7B

HEAVY CHAIN V REGION MODULE GENE SYNTHESIS





FIG. 8






FIG. 10




FIG. 11

pING2006E MET Gly Trp Ser Tyr Ile Ile Leu Phe Leu Val Ala Thr Ala Arg Asp
 GGA TCC₄CCC ACC ATG GGA TGG AGC TAT ATC CTC TTT TTG GTA GCA ACA GCT ACA GAT
 pING2012E GGA TCT CTC GAC ATG 2 30 45 60
 Val His Ser Gln Val Gln Leu Gln Gln Pro Gly Ala Glu Leu Val Lys Pro Gly Ala Ser
 CTC CAC TCC CAG CTC CAA TTG CAG CAG CCT GGG GCT GAA CTG GTG GAA CCA CTT GGG GCT TCA
 75 90 105 120
 Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr Trp MET His Trp Val
 CTG AAG CTG TCC TGC AAG GCT TCT GGC TAC ACC TTC ACC AGC TAC TGG ATG CAC TGG GTC
 135 150 165 180
 Lys Gln Arg Pro Gly Gln Gly Leu Asp Trp Ile Gly Glu Ile Asn Pro Ser Asn Gly Arg
 AAG CAG AGG CCT GGA CAA GGC CTT GAC TGG ATT GGA GAG ATT AAT CCT AGC AAC GGT CGT
 195 210 225 240
 Thr Asn Tyr Asn Glu Lys Phe Lys Ser Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser
 ACT AAT TAC AAT GAG AAG TTC AAG AGC AAG GCC ACA CTG ACT 285 300
 Thr Ala Tyr MET Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys Ala
 ACA GCC TAC ATG CAA CTC AGC AGC CTG ACA TCT GAG GAC TCT CCG GTC TAT TAC TGT
 315 330 345 360
 Ser Tyr Asp Tyr Trp Phe Ala Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 TCC TAT GAT TAC GAC TGG TTT GCT TAC TGG GGC CAA GGC ACT GTG GTC ACC GTC TCC TCA
 375 390 405 420

Bst/EII

FIG.12B




FIG.12A




FIG. 13




FIG. 14

OG ATC CCC CCC CCC CCC CAG TTT GTC TTA AGC CAC TGA CCG CAA GTC TTA GAC ATC ATG GAT TGG CTG TGG AAC TTG 30
 15
 Leu Phe Leu MET Ala Ala Gln Ser Ala Gln Ala Gln Ile Gln Leu Val Gln Ser Gly Pro Gln Leu Lys Lys Pro Gly Gln Thr Val 45
 CTA TTC CTG ATG GCA GCT GGC CAA AGT GGC CAA GCA CAG ATC CAC CAG TCT GCA CAG TCT GAT CAG CTG ANG ANG CAG GCA GCA 180
 165
 Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asn Tyr Gly MET Asn Trp Val Lys Gln Ala Pro Gly Lys Gly Leu Lys Trp MET 150
 135
 ARG ATC TCC TCC AGC GCT TCT GGG TAT ACC TTC ACA AAC TAT GGA ATG AAC TGG CTG ANG CAG GCT CCA GGA ANG GAT TTA ANG TGG 270
 210
 Bel 11 COR2 Lys Trp Ile Asn Thr Tyr Thr Gly Gln Pro Thr Tyr Ala Asp Phe Lys Gly Arg Phe Ala Phe Ser Leu Gln Thr Ser Ala Tyr Thr 360
 315
 GGC TGG ATA MAC ACC TAC ACT GGA CAG CCA ACA TAT GAT CAG TTC ANG GGA CCG TTT GCT TTC TCT TTT GAA ACC TCT GCT TAC ACT 345
 330
 Ala Tyr Leu Gln Ile Asn Asn Leu Lys Asn Gln Asp MET Ala Thr Tyr Phe Cys Ala Arg Phe Ser Tyr Gly Asn Ser Arg Tyr Ser Asp 450
 405
 COR3 Trp Trp Gly Gln Gly Thr Leu Thr Val Ser Ser Ala Lys Thr Ala Pro Ser Val Tyr Pro Leu Ala Pro Val Oys Gly Asp Thr 510
 480
 TAC TGG GGC CAA GGC ACC ACT CTC ACA GTC TCC TCA CCG AAA ACA ACA CCA CCG CCA TCG GTC TAT CCA CTG GCT CCT GAT TGT GGA GAT 540
 465
 Thr Gly Ser Ser Val Thr Leu Gly Cys Leu Val Lys Gly Tyr Phe Pro Gln Pro Val Thr Leu Thr Trp Asn Ser Gly 625
 555
 ACT GGC TCC TGG GTC ACT CTA GGA TGC CTG GTC CAG GGT TAT TTC CCT GAG CCA GTC ATG TTG ACC TGG AAC TCT GGA 600
 585






FIG. 15



BenchI
 16VH CG ATC CCC CCC CCC CCC CCC CAG TTT CTC TTA AGG CAC CAC TGA CCC CAA CTC TTA GAC ATC ATG GAT TCG CTG TCG AAC TTG 62
 PH3-60
 13
 MET Asp Tyr Leu Tyr Asn Leu
 CI-44
 SolI
 CTC GAC TCT AGG CAC CAC TGA CCC CAA CTC TTA GAC ATC ATG GAT TCG CTG TCG AAC TTG
 CI-421
 SolI
 GT CGA CTC TAC TTT GTC TTA AGG CAC CAC TGA CCC CAA CTC TTA GAC ATC ATG GAT TCG CTG TCG AAC TTG

FIG. 17a








FIG.18

A. 5' L6V_K

B.




FIG. 19




FIG.20

2H7 heavy chain variable sequence

leader

C₃₃ GTACCTCTCTACACTCCCTGAAGACACTGACTCTAACCATG GGA TTC AGC AGG ATC TTT

met gly phe ser arg ile phe
NcoI FRI

leu phe leu leu ser val thr thr gly val his ser¹ gln ala tyr leu gln
CTC TTC CTC CTG TCA GTA ACT ACA GGT GTC CAC TCC CAG GCT TAT CTA CAG

gln ser gly ala gln leu val arg pro gly ser val lys met ser cys
CAG TCT GGG GCT GAG CTG GTG AGC CCT GGG GCC TCA GTG AAG ATG TCC TCG

FRI CDR1 CDR1 FR2

lys ala ser gly tyr thr phe thr ser tyr asn met his trp val lys gln
AAG GCT TCT GGC TAC ACA TTT ACC AGT TAC AAT ATG CAC TGG GTA AAG CAG

FR2 CDR2

thr pro Arg gln gly leu glu trp ile gly ala ile tyr pro gly asn gly
ACA CCT AGA CAG GGC CTG GAA TGG ATT GGA GCT ATT TAT CCA GGA AAT GGT

CDR2 FR3

asp thr ser tyr asn gln lys phe lys gln thr leu thr val asp
GAT ACT TCC TAT AAT CAG AAG TTC AAG GGC AAG GCC ACA CTG ACT GTA GAC

lys ser ser ser thr ala tyr met gln leu ser ser leu thr ser glu asp
AAA TCC TCC AGC ACA GCC TAC ATG CAG CTC AGC AGC CTG ACA TCT GAA GAC

FR3 CDR3

ser ala val tyr phe cys ala arg val val tyr tyr ser asn ser tyr trp
TCT GCG GTC TAT TCT TGT GCA AAG GTG GTG TAC TAT AGT AAC TCT TAC TGG

CDR3 FR4

tyr phe asp val trp gln thr gly thr thr val thr val er² FR4
TAC TTC GAT GTC TGG GGC ACA GGG ACC ACG GTC ACC GTC TCG₃₀

↓

Bst EI

J_B Bst EI primer

FIG. 21

2H7 light chain variable sequence

```

met asp phe gln val gln ile phe ser phe leu leu
C23 CCCCAAATTCCAAGACAAATG GAT TTT CAA GTG CAG ATT TTC AGC TTC CTG CTA
      GTC Sali primer FRI
ile ser ala ser val ile ile ala arg gly1 gln ile val leu ser gln ser
ATC AGT GCT TCA GTC ATA ATT GCC AGA GGA CAA ATT GTT CTC TCC CAG TCT
      FRI
pro ala ile leu ser ser pro gln gly lys val thr met thr cys arg1
CCA GCA ATC CTG TCT GCA TCT CCA GGG GAG AGC GTC ATA ATG ACT TCG AGC
CDR1 CDR1, FR2
ala ser ser ser val ser tyr met his1 trp1 tyr gln gln lys pro gly ser
GCC AGC TCA AGT GTA AGT TAC ATG CAC TGG TAC CAG CAG AAG CCA GGA TCC
      KpnI CDR2, FR3 BamHI
ser pro lys pro trp ile tyr1 ala pro ser asn leu ala ser1 gly val pro
TCC CCC AAA CCC TGG ATT TAT GCC CCA TCC AAC CTG GCT TCT GGA GTC CCT
      FR3, CDR3
act arg phe ser gln arg ser gln thr ser tyr ser leu thr ile ser
GCT CGC TTC AGT GGC AGT GGG TCT GGG ACC TCT TAC TCT CTC ACA ATC AGC
      FR3, CDR3
arg val glu ala glu asp ala ala thr tyr tyr cys1 gln gln trp ser phe
AGA GTG GAG GCT GAA GAT GCT GCC ACT TAT TAC TGC CAG CAG TGG AGT TTT
      CDR3, FR4 K FR4
asn pro pro thr phe gly ala gly thr lys leu gln lys
ACA CCA CCC ACG TTC GGT GCT GGG ACC AAG CTG GAG CTG AAA
      T

```

FIG. 22




FIG. 23

2H7 Chimerae




FIG. 24




FIG.25A




FIG. 25B




FIG. 25C




FIG. 26A




FIG. 26 B




FIG.26C




FIG. 27A




FIG.28




FIG. 30




FIG. 27B




FIG. 29




FIG. 31




FIG. 32A




FIG. 33A





FIG. 32B

**FIG. 33B**




FIG. 34

DraI

TTT AAA AGG AAA TTT TTT CTT ATA AAA

CCC AAA TTA TCC AAT CAT CAG TAT TAC AAA ATG TTT CAA CCG TAA TAC ATT TAA CAT TTC

ACC CTT GAA CTG ATC TTA TTT TTT GAC CAC ACT CCC CTT GGT TTT TCA CCA AAA CTG AGT

NdeI

TTC ATT TTT GTT GAA AAA TTT GTA CCT GCG ACA TCG GCG ATA TCG AAC CAT AAA TGC CCA

1

TGA AAA TTC TAT TTC AAG GAG ACA GTC ATA MET Lys Tyr Leu Leu Pro Thr Ala Ala Ala
ATG AAA TAC CTA TTG CCT ACG GCA GCC GCT

HaeIII

Gly Leu Leu Leu Leu Ala Ala Gln Pro Ala MET Ala Ala Asn Thr Gly Gly Tyr Ala Thr
GGA TTC TTA TTA CTC GCT GCC CAA CCA GCG ATG GCC GCA AAT ACG GGT GGC TAT GCC ACC

↑

FIG. 36 A




FIG. 35




FIG. 37A




FIG. 36B




FIG. 38A




FIG. 37B




FIG. 38B




FIG. 39




FIG. 40

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.